


CLAIMS

WE CLAIM

1. A method comprising reacting

- 5 a) at least one diacidic monomer, comprising about 1 to 100 mole % of at least one light-absorbing monomer having a light absorption maximum between about 300 nm and about 1200 nm and 99-0 mole % of a non-light absorbing monomer which does not absorb significant light
10 at wavelengths above 300 nm or has a light absorption maximum below 300 nm, with
b) an organic compound having the formula




15

SUB
C17

- wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene- C₃- C₈-cycloalkylene- C₁-C₄ alkylene, C₁-C₄ alkylene-arylene- C₁-C₄ alkylene, C₂- C₄-alkylene-L-
20 arylene-L- C₂-C₄ alkylene and C₂-C₄ alkylene-(L- C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from the group consisting of -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; X and X₁ are
25 reactive groups and are independently selected from the group consisting of bromine, iodine and R-SO₂O; wherein R is selected from the group consisting of C₁-C₆ alkyl; C₁-C₆ alkyl substituted with chlorine, fluorine, C₁-C₆ alkoxy, aryl, aryloxy, arylthio or C₃-C₈ cycloalkyl; C₃-C₈
30 cycloalkyl or aryl,
wherein said reaction is carried out in a solvent in the presence of a base to form a light absorbing polymeric composition having the formula

09751766-122900



wherein B is as defined above, n is at least 2 and A comprises the residue of said diacidic monomer.

5

2. The process of claim 1 where said light-absorbing monomers have the formula



10

wherein H represents an acidic hydrogen atom; Y is a divalent light-absorbing moiety selected from the group consisting of chromophoric classes of azo, disazo, triazo, bis-azo, methine, arylidene, polymethine, azo-methine, azamethine, anthraquinone, anthrapyridone (3H-dibenz[f,ij]isoquinoline-2,7-dione, nitroarylamines anthrapyridine (7H-dibenz[f,ij]isoquinoline-7-one, phthaloylphenothiazine (14H-naphth[2,3-a]phenothiazine-8,13-dione, benzanthrone (7H(de)anthracene-7-one), anthrapyrimidine (7H-benzo[e]perimidine-7-one), anthrapyrazole, anthraisothiazole, triphenodioxazine, thiaxanthene-9-one, fluorindine (5,12-dihydroquinoxaline [2,3-b]phenazine, quinophthalone, phthalocyanine, metal phthalocyanine, naphthalocyanine, metal naphthalocyanine, nickel dithiolenes, squarylium compounds, croconium compounds, coumarin (2H-1-benzopyran-2-one), coumarin imine (2H-1-benzopyran-2-imine), perinone, benzodifuran, phthaloylacridone, phthaloylphenoxazine (14H-naphtho[2,3-a]phenoxazine-8,13-dione, phthaloylacridone (13H-naphtho[2,3-c] acridine-5,8,14-trione), anthraquinonethioxanthene (8H-naphtho[2,3-c]thioxanthene-5,8,13-trione, anthrapyridazone, pyrrolo[3,4-c]pyrrole, indigo, thioindigo, quinoline, xanthene, acridine, azine, cyanine, oxazine, 1,4 and 1,5-naphthoquinones, 2,5-

SUB
C17

009751766-122900

09751766-12900

diarylaminoterephthalic acids and esters, pyromellitic acid diimide, naphthalene-1,4,5,8-tetracarboxylic acid diimide, 3,4,9,10-perylenetetracarboxylic acid diimide, 3-aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-oxopyrroline, arylisoindoline, hydroxybenzophenone, benoztriazole, naphthotriazole, diminoisoindoline, naphthopyran (3H-naphtho[2,1-6]pyran-3-one and 3-imine, phthalimides, 2-arylbenzazoles, carbostyryls, 1,2-diarylethenes, 2,5-diarylthiophenes, 2,5-diaryl-1,3,4-oxadiazoles, triazines, 2,5-diarylfurans, 2,5-diaryl-1,3,4-thiadiazoles, thiophenes, 1,3-diphenyl-2-pyrazolines, 2-arylbenzofurans, 2,6-diphenylbenzofurans, quinolines, quinoxalines, 3,4-diarylfuanones, distyrylarenes, benzanthrones, polyarenes and naphthalimides.


SUB
C7

3. The method of claim 2 wherein the hydrogen atoms of said formula H-Y-H are independently bonded to an oxygen, sulfur, or nitrogen atom which is a part of the light absorbing moiety Y and which in combination provides two acidic functional groups.

4. The method of claim 3 wherein said acidic functional groups have pKa values of from about 1.5 to about 12.

5. The method of claim 3 wherein said acidic functional groups are independently selected from the group consisting of -CO₂H, -SH, -OH attached to an aromatic ring, -CONHCO-, -SO₂-NH-CO-, -SO₂-NH-SO₂-, 1(H)-1,2,4-triazol-3-yl-, imidazolyl, benzimidazolyl, pyrazolyl, -SO₂H attached to aromatic ring, -NHSO₂R₅ and -SO₂NHR₅, wherein R₅ is selected from the group consisting of C₁-C₆ alkyl, C₃-C₈ cycloalkyl, aryl and C₁-C₆ alkyl substituted with at least one group selected from the group consisting of C₁-C₆ alkoxy, aryl, aryloxy, arylthio and C₃-C₈ cycloalkyl.

6. The method of claim 1 wherein said non light-absorbing monomers have the formula




wherein H represents an acidic hydrogen atom; Y₁ is a
 5 divalent moiety selected from the group consisting of -O₂C-
 R₁-CO₂- and -O-R₂-O- and -O₂C-R₃-O-, wherein R₁ is selected
 from the group consisting of C₂-C₁₂ alkylene, 1-4-
 cyclohexylene, arylene, arylene-O-arylene, arylene-SO₂-
 arylene, arylene-S-arylene, and C₁-C₄ alkylene-O- C₁-C₄
 10 alkylene; wherein R₂ is selected from the group consisting
 of arylene, arylene-O-arylene, arylene-S-arylene, arylene-
 SO₂-arylene, phenylene-phenylene, and phenylene-C(R₄)₂-
 phenylene; wherein R₄ is selected from the group
 consisting of hydrogen and C₁-C₄ alkyl; wherein R₃ is
 15 selected from arylene.

SUB
C17

7. The method of claim 1 wherein said polymeric
 composition is linear.

8. The method of claim 1 wherein said diacidic
 monomers have pK_a values of about 12 or below.

20 9. The method of claim 2 wherein H-Y-H includes a
 moiety selected from the group consisting of carboxy
 groups attached to an aromatic ring carbon or aliphatic
 carbon, hydroxy groups attached to an unsubstituted or
 substituted phenyl or naphthyl radical, -CO-NHCO- groups
 25 attached to an aromatic ring to provide an imide and
 1(H)-1,2,4-triazol-3-yl group having the formula



30 wherein R₅' is selected from the group consisting of
 hydrogen, C₁-C₆ alkyl and aryl.

10. The method of claim 1 where n is between about
 2 and about 25.

005231-994560

11. The method of claim 1 wherein n is between about 3 and about 15.

SUB
C17

5

12. The method of claim 1 wherein said base is selected from the group consisting of alkali metal carbonates, alkali metal bicarbonates and tertiary amines, aromatic nitrogen bases, bicyclic nitrogen containing bases having non-hindered electron pairs and mixtures thereof.

Sub B1

10

13. The method of claim 12 wherein said base is selected from the group consisting of triethylamine, tri-n-butylamine, N-methylpiperidine, N,N'-dimethylpiperazine, N-methylmorpholine and N,N,N',N'-tetramethylethylenediamine, pyridines, picolines, quinolines, isoquinolines, N-alkylpyrroles, N-alkylimidazoles, 1,8-diazabicyclo[5,4,0]undec-7-ene (DBU), 1,5-diazabicyclo[4,3,0]non-5-ene (DBN) and 1,4-diazadicyclo[2,2,2]octane (DABCO®) and mixtures thereof.

15

14. The method of claim 1 wherein said solvent is one or more aprotic polar solvents.

20

15. The method of claim 1 wherein said solvent is selected from the group consisting of N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidone, N-methyl-N-phenyl formamide, dimethylsulfoxide, aliphatic nitriles, sulfolane, hexamethyl phosphoramidate, water, alcohols, ketones pyridine and ether-alcohols and mixtures thereof.

SUB

25

C17

16. The method of claim 15 wherein said solvent is selected from the group consisting of N,N-dimethylacetamide, N,N-dimethylformamide, N-methyl-2-pyrrolidone, N-methyl-N-phenyl formamide, dimethylsulfoxide, aliphatic nitriles, sulfolane and hexamethyl phosphoramidate and mixtures thereof.


30

17. The method of claim 1 wherein said reacting is conducted at a temperature between about 75°C and about 125°C.

35

0054756-122900

18. The method of claim 1 wherein said organic compound having the formula



5

SUB
C17

is selected from the group consisting of disulfonate compounds where X and X₁ are both a sulfonate ester of the formula-OSO₂R, wherein R is selected from C₁-C₄ alkyl, phenyl or p-methylphenyl and wherein B is selected from C₂-C₆ alkylene, -CH₂-1,4-cyclohexylene-CH₂-, -CH₂CH₂(OCH₂CH₂)₁₋₄ and-CH₂CH₂O-1,4-phenylene-O-CH₂CH₂-.

10

SUB
C17

19. The method of claim 18 wherein said B moiety of organic compound of Formula II is selected from the group consisting of-CH₂CH₂-, -CH₂CH(CH₃)CH₂-, -CH₂C(CH₃)₂CH₂-, -(CH₂)₄-, -(CH₂)₆-, -CH₂CH₂OCH₂CH₂- and-CH₂-1,4-cyclohexylene-CH₂-.

15

20. The method of claim 1 wherein said organic compound having the formula X-B-X₁ is selected from the group consisting of 1,2-ethandiol, dimethanesulfonate; 1,2-ethanediol bis(4-methylbenzenesulfonate); 1,4-butanediol, dimethane sulfonate; 1,6-hexanediol, dimethanesulfonate; 1,3-propanediol,2,2-dimethyl-, dimethanesulfonate; 1,4-cyclohexanedimethanol, dimethanesulfonate; 1,1,3,3-tetramethylcyclobutanediol, dimethanesulfonate, and ethanol 2,2'-oxybis-dimethanesulfonate.

20

25

SUB
C17


21. The method of claim 1 wherein A of said light absorbing polymeric composition comprises 100 mole% of said light-absorbing monomer.

30

22. The method of claim 1 wherein said diacidic light absorbing monomer has a light absorption maximum between about 300 nm and about 1200 nm and is present in said light absorbing polymeric composition in an amount at least about 50% by weight %.

35

23. A light absorbing composition having the formula



wherein A₁ comprises the residue of at least one diacidic monomer having a light absorption maximum between about 300 nm and about 1200 nm, and wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene- C₃-C₈-cycloalkylene- C₁-C₄ alkylene, C₁-C₄ alkylene-arylene- C₁-C₄ alkylene, and C₂- C₄-alkylene-L-arylene-L- C₂-C₄ alkylene and C₂-C₄ alkylene-(L- C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from-O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; wherein n is at least 2.

24. A composition comprising a thermoplastic polymer blended with at least one light absorbing linear polymeric composition of claim 23.

25. The composition of claim 24 wherein said thermoplastic polymer is selected from the group consisting of polyesters, polyolefins, polyamides, polyimides, polyvinyl chloride, polyurethanes, polycarbonates, cellulose esters, polyacrylates, polyvinylesters, polyester-amides, polystyrene, polyacrylonitrile-butadiene- styrene, polystyrene-acrylonitrile and mixtures and blends thereof.

26. The composition of claim 23 wherein A₁ comprises the residue of at least one diacidic monomer having the structure



wherein H represents an acidic hydrogen atom and Y is a divalent light-absorbing moiety selected from the chromophoric classes consisting of azo, disazo, triazo, bis-azo, methine, arylidene, polymethine, azo-methine, azamethine, anthraquinone, anthrapyridone (3H-dibenz[f,i,j] isoquinoline-2,7-dione, nitroarylamines anthrapyridine

09751766-122900


SUB
C7

00622T 997560

(7H-dibenz[f,i]isoquinoline-7-one, phthaloylphenothiazine
 (14H-naphth[2,3-a]phenothiazine-8,13-dione, benzanthrone
 (7H(de)anthracene-7-one), anthrapyrimidine (7H-
 benzo[e]perimidine-7-one), anthrapyrazole,
 5 anthraisothiazole, triphenodioxazine, thiaxanthene-9-one,
 fluorindine (5,12-dihydroquinoxaline [2,3-b]phenazine,
 quinophthalone, phthalocyanine, metal phthalocyanine,
 naphthalocyanine, metal naphthalocyanine, nickel
 dithiolenes, squarylium compounds, croconium compounds,
 10 coumarin (2H-1-benzopyran-2-one), coumarin imine (2H-1-
 benzopyran-2-imine), perinone, benzodifuran,
 phthaloylacridone, phthaloylphenoxazine (14H-naphtho[2,3-
 a]phenoxazine-8,13-dione, phthaloylacridone (13H-
 naphtho[2,3-c]acridine-5,8,14-trione),
 SUB 15 anthraquinonethioxanthene (8H-naphtho[2,3-c]thioxanthene-
 C17 5,8,13-trione, anthrapyridazone, pyrrolo[3,4-c]pyrrole,
 indigo, thioindigo, quinoline, xanthene, acridine, azine,
 cyanines, oxazine, 1,4 and 1,5-naphthoquinones, 2,5-
 diarylaminoterephthalic acids and esters, pyromellitic
 20 acid diimide, naphthalene-1,4,5,8-tetracarboxylic acid
 diimide, 3,4,9,10-perylenetetracarboxylic acid diimide, 3-
 aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-
 oxopyrroline, arylisoindoline, hydroxybenzophenone,
 benotriazole, naphthotriazole, diminoisoindoline,
 25 naphthopyran (3H-naphtho[2,1-6]pyran-3-one and 3-imine,
 phthalimides, 2-arylbenzazoles, carbostyryls, 1,2-
 diarylethenes, 2,5-diarylthiophenes, 2,5-diaryl-1,3,4-
 oxadiazoles, triazines, 2,5-diarylfurans, 2,5-diaryl-
 1,3,4-thiadiazoles, thiophenes, 1,3-diphenyl-2-
 30 pyrazolines, 2-arylbenzofurans, 2,6-diphenylbenzofurans,
 quinolines, quinoxalines, 3,4-diarylfuanones,
 distyrylarenes, benzanthrone, polyarenes and
 naphthalimides.

27. The light absorbing linear polymeric
 35 composition of claim 23 or 26

wherein A₁ further comprises less than about 50% by weight of the total composition of a residue of at least one non-light absorbing monomer having the formula



5

wherein Y₁ is a divalent moiety, selected from the group consisting of -O₂C-R₁-CO₂- and -O-R₂-O- and -O₂C-R₃-O-, wherein

10

R₁ is selected from the group consisting of C₂-C₁₂ alkylene, 1-4-cyclohexylene, arylene, arylene-O-arylene, arylene-SO₂-arylene, arylene-S-arylene, and C₁-C₄ alkylene-

15

O- C₁-C₄ alkylene; wherein R₂ is selected from the group consisting of arylene, arylene-O-arylene, arylene-S-arylene, arylene-SO₂-arylene, phenylene-phenylene, and phenylene-C(R₄)₂-phenylene; wherein R₄ is selected from the group consisting of hydrogen and C₁-C₄ alkyl; wherein R₃ is

SUB
C17

20

arylene; wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene-C₃-C₈-cycloalkylene-C₁-C₄ alkylene, C₁-C₄ alkylene-arylene-C₁-C₄ alkylene, and C₂-C₄-alkylene-L-arylene-L-C₂-C₄ alkylene and C₂-C₄ alkylene-(L-C₂-C₄ alkylene)₁₋₄; wherein L is a linking group selected from -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂ aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; wherein n is at least 2.

25

28. The light absorbing linear polymeric composition of Claim 25 wherein A₁ comprises the residue of at least one diacidic monomer having at least one acidic group other than carboxy and having a light absorption maximum between about 300 nm and about 1200 nm

30

and wherein B is a divalent organic radical selected from C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene-C₃-C₈-cycloalkylene-C₁-C₄ alkylene, C₁-C₄ alkylene-arylene-C₁-C₄ alkylene, and C₂-C₄-alkylene-L-arylene-L-C₂-C₄ alkylene and C₂-C₄ alkylene-(L-C₂-C₄ alkylene)₁₋₄; wherein L

35

is a linking group selected from -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-,

09751766.122900

-(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; wherein n is at least 2.

29. The process of claim 2 wherein said light absorbing monomer comprises two carboxy groups as acidic functional groups.

30. The process of claim 2 wherein said light absorbing monomer comprises two 1(H)-1,2,4-triazol-3-ylthio groups as acidic functional groups.

31. The process of claim 2 wherein said light absorbing monomer comprises one carboxy and one 1(H)-1,2,4-triazol-3-ylthio group as acidic functional groups.

32. The process of claim 2 wherein said light absorbing monomer comprises two hydroxy groups attached to aromatic ring(s).

33. The process of claim 2 wherein said light absorbing monomer comprises one imide group and one carboxy group.

34. The process of claim 2 wherein said light absorbing monomer comprises one imide group and one 1(H)-1,2,4-triazol-3-ylthio group.

35. The composition of claim 25 wherein said light absorbing monomer comprises two carboxy groups as acidic functional groups.

36. The composition of claim 25 wherein said light absorbing monomer comprises two 1(H)-1,2,4-triazol-3-ylthio groups as acidic functional groups.

37. The composition of claim 25 wherein said light absorbing monomer comprises one carboxy and one 1(H)-1,2,4-triazol-3-ylthio group as acidic functional groups.

38. The composition of claim 25 wherein said light absorbing monomer comprises two hydroxy groups attached to aromatic ring(s).

39. The composition of claim 25 wherein said light absorbing monomer comprises one imide group and one carboxy group.

005221 997560

SUB
C17

40. The composition of claim 23 wherein said light absorbing monomer comprises one imide group and one 1(H)-1,2,4-triazol-3-ylthio group.

41. The composition of claim 23 wherein said light absorbing monomer comprises a diacidic sulfamoyl ($-\text{SO}_2\text{NH}_2$) group.

42. The composition of claim 25 wherein said light absorbing monomer comprises two acidic groups independently selected from the group consisting of $-\text{CO}_2\text{H}$, SH, hydroxy attached to an aromatic ring, $-\text{CONHCO}$ - (imide), $-\text{SO}_2\text{NHCO}$ -, $-\text{SO}_2\text{NHSO}_2$ -, 1(H)-1,2,4-triazolyl-3-yl-, imidazolyl, benzimidazolyl, pyrazolyl, $-\text{SO}_2\text{H}$ attached to an aromatic ring, $-\text{NHSO}_2\text{R}_5$ and $-\text{SO}_2\text{NHR}_5$, wherein R_5 is selected from the group consisting of C_1 - C_6 alkyl; C_1 - C_6 alkyl substituted with at least one group selected from C_1 - C_6 alkoxy, aryl, aryloxy, arylthio and C_3 - C_8 cycloalkyl; C_3 - C_8 cycloalkyl; aryl.

43. The composition of claim 27 wherein said light absorbing monomer comprises two carboxy groups as acidic functional groups.

44. The composition of claim 27 wherein said light absorbing monomer comprises two 1(H)-1,2,4-triazol-3-ylthio groups as acidic functional groups.

45. The composition of claim 27 wherein said light absorbing monomer comprises one carboxy and one 1(H)-1,2,4-triazol-3-ylthio group as acidic functional groups.

46. The composition of claim 27 wherein said light absorbing monomer comprises two hydroxy groups attached to aromatic ring(s).

47. The composition of claim 27 wherein said light absorbing monomer comprises one imide group and one carboxy group.

48. The composition of claim 27 wherein said light absorbing monomer comprises one imide group and one 1(H)-1,2,4-triazol-3-ylthio group.

09751766-122900

SUB
C17

49. The composition of claim 27 wherein said light absorbing monomer comprises a diacidic sulfamoyl ($-\text{SO}_2\text{NH}_2$) group.

50. The composition of claim 27 wherein said light
5 absorbing monomer comprises two acidic groups independently selected from $-\text{CO}_2\text{H}$, SH, hydroxy attached to an aromatic ring, $-\text{CONHCO}-$ (imide), $-\text{SO}_2\text{NHCO}-$, $-\text{SO}_2\text{NHSO}_2-$, 1(H)-1,2,4-triazolyl-3-yl-, imidazolyl, benzimidazolyl, pyrazolyl, $-\text{SO}_2\text{H}$ attached to an aromatic ring, $-\text{NHSO}_2\text{R}_5$
10 and $-\text{SO}_2\text{NHR}_5$, wherein R_5 is selected from C_1 - C_6 alkyl; C_1 - C_6 alkyl substituted with at least one group selected from C_1 - C_6 alkoxy, aryl, aryloxy, arylthio and C_3 - C_8 cycloalkyl; C_3 - C_8 cycloalkyl; aryl.

51. The light absorbing linear polymeric composition
15 of claim 27 wherein said at least one diacidic monomer comprises at least about 50% by weight of the total composition.









52. The composition of claim 51 wherein the light
absorbing portion of A comprises the residue of at least
20 one diacidic light absorbing monomer selected from the group consisting of the anthraquinone and anthrapyridone colorants having the structures:






09751766-122900

SUB
C17

006221-9975/60

SUB
C1





SUB
C7

wherein R_{14} is selected from the group consisting of hydrogen and 1-4 groups selected from amino, C_1 - C_{10}


5 alkylamino, C_3 - C_8 alkenylamino, C_3 - C_8 alkynylamino, C_3 - C_8 cycloalkylamino, arylamino, halogen, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, aryl, aroyl, C_1 - C_6 alkanoyl, C_1 - C_6 alkanoyloxy,

$NHCO$ C_1 - C_6 alkyl, $NHCO$ aryl, $NHCO_2$ C_1 - C_6 alkyl, $NHSO_2$ C_1 - C_6 alkyl, $NHSO_2$ aryl, C_1 - C_6 alkoxycarbonyl, aryloxy, arylthio,

10 heteroarylthio, cyano, nitro, trifluoromethyl, thiocyno, SO_2 C_1 - C_6 alkyl, SO_2 aryl, $-SO_2NH$ C_1 - C_6 alkyl, $-SO_2N(C_1$ - C_6 alkyl) $_2$, $-SO_2N(C_1$ - C_6 alkyl) aryl, $CONH$ C_1 - C_6 alkyl, $CON(C_1$ - C_6 alkyl) $_2$, $CON(C_1$ - C_6 alkyl) aryl, C_1 - C_6 alkyl, furfurylamino, tetrahydrofurfurylamino, 4-(hydroxymethyl)


15 cyclohexanemethylamino,

09751766-122900



or hydroxy; Q and Q' are independently selected from the
 5 group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-,
 -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is
 selected from the group consisting of hydrogen, aryl, C₃-C₈
 cycloalkyl, or C₁-C₁₀ alkyl; R₁₅ is selected from the group
 10 consisting of hydrogen, cyano, C₁-C₆ alkylamino, C₁-C₆
 alkoxy, halogen, arylthio, aryl, heteroaryl,
 heteroarylthio, C₁-C₆ alkoxy carbonyl, aroyl or
 arylsulfonyl; R₁₆ is selected from the group consisting of
 hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl and aryl; R₁₆' is
 selected from the group consisting of hydrogen, C₁-C₆
 15 alkyl, halogen and C₁-C₆ alkoxy; wherein each C₁-C₆ alkyl
 group and C₁-C₆ alkyl group which is a portion of another
 group may contain at least one substituent selected from
 the group consisting of hydroxy, cyano, chlorine,
 fluorine, C₁-C₆ alkoxy, C₃-C₈ cycloalkoxy, C₁-C₆
 20 alkylcyclohexyl, hydroxymethyl cyclohexyl, aryl and
 heteroaryl; with the provision that two acidic groups
 containing one acidic proton each or one acidic group
 containing two acidic hydrogens be present in the diacidic
 compounds.

25 53. The composition of claim 26 or 27 wherein the
 light absorbing portion of A comprises the residue of at
 least one light absorbing monomer selected from the group
 consisting of azo, disazo, bis-azo and azomethine and
 having respectively the structures:



wherein R₆ is the residue of an aromatic or heteroaromatic
 amine which has been diazotized and coupled with a
 35 coupling component H-Z and is derived from an amine

09751766 133500

SUB
C17

Sub B3

Sub B
cont.


09751766-122900

- selected from the group consisting of aromatic and heteroaromatic amine classes selected from the group consisting of aniline, 1-aminonaphthalene, 1-aminoanthraquinone, 4-aminoazobenzene, 2-aminothiazole, 2-aminobenzothiazole, 3-amino-2,1-benzisothiazole, 2-aminothieno[2,3-d]thiazole, 5-aminoisothiazole, 5-aminopyrazole, 4-aminopyrazoloisothiazole, 2-amino-1,3,4-thiadiazole, 5-amino-1,2,4-thiadiazole, 5-amino-1,2,3-triazole, 2-amino-1,3,4-triazole, 2(5) aminoimidazole, 3-aminopyridine, 2(3) aminothiophene, 2(3) aminobenzo[b]thiophene, 2-aminothieno[3,2-b]thiophene, 3-aminothieno[2,3-c]isothiazole, 3-amino-7-benz-2,1-isothiazole, 3-aminobenzothienoisothiazole, 3-aminoisothiazole[3,4-d]pyrimidine, 5-amino-1,2,3-triazole, 3(4) aminophthalimide and 5(6) amino-1,2-benzisothiazolon-1,1-dioxide with said aromatic and heteroaromatic ring systems being unsubstituted or substituted with one or more groups selected from C₁-C₁₀ alkyl, C₁-C₆ alkoxy, C₃-C₈ cycloalkyl, carboxy, halogen, C₁-C₆ alkoxycarbonyl, formyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy, dicyanovinyl, C₃-C₈-cycloalkanoyl, thiocyno, trifluoroacetyl, cyano, carbamoyl, -CONH-C₁-C₆ alkyl, CONHaryl, CON(C₁-C₆ alkyl)₂, sulfamoyl, SO₂NH C₁-C₆ alkyl, SO₂N(C₁-C₆ alkyl)₂, SO₂NHaryl, SO₂NH C₃-C₈ cycloalkyl, CONH C₃-C₈ cycloalkyl, aryl, aroyl, -NHSO₂ C₁-C₆ alkyl, -N(C₁-C₆ alkyl)SO₂ C₁-C₆ alkyl, -NHSO₂ aryl, NHCO C₁-C₆ alkyl, NHCO C₃-C₈ cycloalkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHCONH C₁-C₆ alkyl, NHCONHaryl, N(C₁-C₆ alkyl)aryl, arylazo, heteroaryl, aryloxy, arylthio, C₃-C₈ cycloalkoxy, heteroarylazo, heteroarylthio, arylsulfonyl, tricyanovinyl, aryloxysulfonyl, C₁-C₆ alkylsulfonyl, trifluoromethyl, fluorosulfonyl, trifluoromethylsulfonyl, thiocyno, hydroxy, nitro or CH=D, wherein D is the residue of an active methylene compound selected from the group consisting of malononitrile, α-cyanoacetic acid esters, malonic acid esters, α-cyanacetic acid amides, α-C₁-C₆

Sub B3
Cont.

00751766-122900

- alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles, α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles, α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane dinitrile-5,5-dioxides, 1,3-bis(dicyanomethylene) indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides or aryl-
- 10 C(CH₃)C=C(CN)₂; wherein R₇ is a divalent aromatic or heteroaromatic radical selected from the group consisting of 1,4-phenylene, naphthalene-1,4-diyl, thiazol-2,5-diyl and thiophene-2,5-diyl:



15

- wherein R₈ is selected from the group consisting of hydrogen or 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆ alkoxy, cyano, halogen, -NHCO C₁-C₆ alkyl, -NHCO₂ C₁-C₆ alkyl, -NHCO aryl, -NHCONH aryl or NHCONH C₁-C₆ alkyl; R₉ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, halogen, aryl, heteroaryl; R₁₀ is selected from the group consisting of hydrogen, C₁-C₆ alkoxycarbonyl, cyano, carbamoyl, aryl, arylsulfonyl, aroyl, -CONH C₁-C₆ alkyl, or C₁-C₆ alkylsulfonyl; wherein Z is the residue of a coupling compound selected from the group consisting of electron rich compound classes of anilines, 1-aminonaphthalenes, 1,2-dihydroquinolines, 1,2,3,4-tetrahydroquinolines, benzomorpholines (3,4-dihydro-2H-1,4-benzoxazine),
- 20
- 25


pyrazolones, pyrazoles, 3-cyano-6-hydroxy-2-pyridones,
2,3-dihydroindoles, indoles, 4-hydroxycoumarins,
4-hydroxy-2-quinolones, imidazo[2,1-b]thiazoles,
julolidines (2,3,6,7-tetrahydro-1H,5H-
5 benzo[ij]quinolizines), 1-oxajulolidines, 1,2,5,6-
tetrahydro-4H-pyrrolo[3,2,1-ij]quinolines, 2,6-diamino-3
cyanopyridines, 2-aminothiazoles, 2-aminothiophenes, 5,5-
dimethyl-1,3-cyclohexanedione (dimedone), phenols,
naphthols, 2,4-pentanediones or acetoacetarilides; wherein
10 Y₁ is the residue of a bis coupling component selected
from the group consisting of anilines, 1,2-
dihydroquinolines, 1,2,3,4-tetrahydroquinolines,
benzomorpholines (3,4-dihydro- 2H-1,4-benzoxazines), 3-
cyano-6-hydroxy-2-pyridones, 2,6-diaminopyridines, 2,3-
15 dihydroindoles, naphthylamines, 2-aminothiazoles, or a
combination of these; with the provision that two acidic
functional groups containing one acidic hydrogen each or a
functional group containing two acidic hydrogen are
present in the diacidic light absorbing monomer.

20 54. The composition of claim 53 wherein Z is
selected from the group consisting of:


Sub B3
cont.

00622T" 99475260

SUB
C/




wherein R_{17} is selected from the group consisting of hydrogen, 1-2 groups selected from C_1-C_6 alkyl, C_1-C_6 alkoxy, C_1-C_6 alkylthio, $-O-C_2-C_6$ alkylene-OH, $O-C_2-C_6$ alkylene- C_1-C_6 alkanoyloxy, C_1-C_6 alkylene-OH, C_1-C_6 alkylene- C_1-C_6 alkanoyloxy, halogen, carboxy, C_1-C_6 alkoxycarbonyl, trifluoromethyl, $NHCO_2R_{24}$, $NHCO_2R_{24}$, $NHCON(R_{24})R_{25}$, and $NHSO_2R_{25}$, wherein R_{24} is selected from the group consisting of hydrogen, C_1-C_{10} alkyl, C_3-C_8 cycloalkyl or aryl, R_{25} is selected from the group consisting of C_1-C_{10} alkyl, C_3-C_8 cycloalkyl or aryl wherein each C_1-C_{10} alkyl group in R_{24} and R_{25} may be further substituted with one or more groups selected from the group consisting of C_3-C_8 cycloalkyl, aryl, aryloxy, arylthio, CO_2H , $CO_2C_1-C_6$ alkyl, cyano, hydroxy, succinimido, C_1-C_6 alkoxy,



wherein R_5' is selected from the group consisting of hydrogen, C_1-C_6 alkyl or aryl; R_{16}' is selected from hydrogen or one or two groups selected from C_1-C_6 alkyl, halogen and C_1-C_6 alkoxy; Q is selected from the group consisting of $-O-$, $-N(COR_{10})-$, $-N(R_{10})-$, $-S-$, $-SO_2-$, $-CO_2-$, $-CON(R_{10})$, $-SO_2(R_{10})-$, wherein R_{10} is selected from the group consisting of hydrogen, aryl, C_3-C_8 cycloalkyl or C_1-C_{10} alkyl; R_{18} and R_{19} are independently selected from the group consisting of hydrogen, unsubstituted C_1-C_{10} alkyl, substituted C_1-C_{10} alkyl, C_3-C_8 cycloalkyl, C_3-C_8 alkenyl, C_3-C_8 alkynyl and aryl or R_{18} and R_{19} may be

combined with another element to which they are attached to form a radical Z having the formula



5


wherein Q₂ is selected from the group consisting of a covalent bond, -O-, -S-, -SO₂-, -CO-, -CO₂-, -N-(C₁-C₆ alkyl)-, -N(COC₁-C₆ alkyl)-, -N(SO₂C₁-C₆ alkyl)-, -N(CO aryl)-, or -N(SO₂ aryl); R₂₀, R₂₁ and R₂₂ are independently selected from the group consisting of hydrogen or C₁-C₆ alkyl; R₂₃ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, heteroaryl or aryl.

55. The composition of claim 26 or 51 wherein the light absorbing portion of A comprises the residue of at least one light absorbing monomer selected from the group consisting of methine, arylidene, polymethine, azamethine, 3-aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-oxypyrroline and arylisindoline and having respectively the structures:

20

SUB
U7

006221-9975760




- wherein R_{11} is the residue of an electron rich aromatic compound selected from the group consisting of an aniline,
- 5 1-naphthylamine, 1,2-dihydroquinoline, 1,2,3,4-tetrahydroquinoline, 1,3,3-trimethyl-2-methyleneindole, 1,3-dihydro-2-methylene-1,1,3-trimethyl-2H-benz[e]indole, imidazo [2,1-b] thiazole, benzomorpholine (3,4-dihydro-2H-1,4,benzoxazine), 2,3-dihydroindole, indole, 2-
- 10 aminothiazole, julolidine (2,3,6,7-tetrahydro-1H, 5H- benz [ij] quinolizine, 1-oxajulolidine, 4H-pyrrolo [3,2,1-ij]-quinoline, phenol, naphthol, thiophenol, pyrrole, pyrazole, furan, thiophene, carbazole, phenothiazine or phenoxazine compound; R_{12} is selected from the group
- 15 consisting of hydrogen, C_1-C_{10} alkyl, C_3-C_8 alkenyl, C_3-C_8 -alkynyl, C_3-C_8 cycloalkyl, aryl, $(CH_2CH_2O)_{1-3}$ R_{13} and C_1-C_4 alkylene- C_3-C_8 cycloalkylene, wherein the C_1-C_6 alkyl groups may be substituted by at least one group selected from the group consisting of carboxy, C_1-C_6 carbalkoxy,
- 20 C_1-C_6 alkanoyloxy, cyano, hydroxy, chlorine, fluorine, C_1-C_6 alkoxy, C_3-C_8 cycloalkyl or aryl; R_{13} is selected from the group consisting of hydrogen, C_1-C_6 alkoxy or C_1-C_6 alkanoyloxy; wherein D is the residue of an active


09754766.122900

SUB
C17

methylene compound selected from the group consisting of malononitrile, α -cyanoacetic acid esters, malonic acid esters, α -cyanacetic acid amides, α -C₁-C₆ alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles, α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles, α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane dinitrile-5,5-dioxides, 1,3-bis(dicyanomethylene) indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides or aryl-C(CH₃)C=C(CN)₂, with the proviso that two acidic functional groups containing one acidic hydrogen each, or a functional group containing two acidic hydrogens are present.

56. The composition of claim 55 wherein R₁₁ is selected from the group consisting of the electron rich aromatic residues corresponding to the structures:






SUB
C17


- wherein R_{26} is selected from the group consisting of hydrogen or a group selected from the group consisting of
- 5 C_1 - C_6 alkoxy, CO_2H , C_1 - C_6 alkyl or C_1 - C_6 alkoxy;
 - wherein R_{17} is selected from the group consisting of hydrogen, and 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, $-O$ C_2 - C_6 alkylene-OH, O C_2 - C_6 alkylene- C_1 - C_6 alkanoyloxy, C_1 - C_6 alkylene-OH, C_1 - C_6 alkylene- C_1 - C_6 alkanoyloxy, halogen, carboxy, C_1 - C_6
 - 10 alkoxy, trifluoromethyl, $NHCOR_{24}$, $NHCO_2R_{24}$, $NHCON(R_{24})R_{25}$, and $NHSO_2R_{25}$, wherein R_{24} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl or aryl, R_{25} is selected from the group consisting of C_1 - C_{10}
 - 15 alkyl, C_3 - C_8 cycloalkyl or aryl wherein each C_1 - C_{10} alkyl group in R_{24} and R_{25} may be further substituted with one or more groups selected from the group consisting of C_3 - C_8 cycloalkyl, aryl, aryloxy, arylthio, CO_2H , CO_2 C_1 - C_6 alkyl,

09751766-122500



SUB
C17

- wherein R_5' is selected from the group consisting of hydrogen, C_1 - C_6 alkyl or aryl; R_{16}' is selected from the group consisting of hydrogen, one or two groups selected from C_1 - C_6 alkyl, halogen and C_1 - C_6 alkoxy; Q is selected from the group consisting of $-O-$, $-N(COR_{10})-$, $-N(R_{10})-$, $-S-$, $-SO_2-$, $-CO_2-$, $CON(R_{10})$, $SO_2(R_{10})-$ wherein R_{10} is selected from the group consisting of hydrogen, aryl, C_3 - C_8 cycloalkyl or C_1 - C_{10} alkyl; R_{18} and R_{19} are independently selected from the group consisting of hydrogen, unsubstituted C_1 - C_{10} alkyl, substituted C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl and aryl or R_{18} and R_{19} may be combined with another element to which they are attached to form a radical Z having the formula






- wherein Q_2 is selected from the group consisting of a covalent bond, $-O-$, $-S-$, $-SO_2-$, $-CO-$, $-CO_2-$, $-N-(C_1-C_6$ alkyl)-, $-N(CO C_1-C_6$ alkyl)-, $-N(SO_2 C_1-C_6$ alkyl)-, $-N(CO$ aryl)-, or $-N(SO_2$ aryl); R_{20} , R_{21} and R_{22} are independently selected from the group consisting of hydrogen or C_1 - C_6 alkyl; R_{23} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, heteroaryl or aryl.

09751766-122900

Sub B4


57. The composition of claim 51 wherein the light absorbing portion of A₂ comprises a residue of at least one diacidic coumarin compounds selected from the group consisting of the structures




- wherein Z₃ is selected from the group consisting of cyano, C₁-C₆ alkoxycarbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl, heteroaryl, formyl, aroyl, C₁-C₆ alkanoyl or -CH=D, wherein R₁₇ is selected from the group consisting of hydrogen, 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, -O C₂-C₆ alkylene-OH, O C₂-C₆ alkylene- C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene- C₁-C₆ alkanoyloxy, halogen, carboxy, C₁-C₆ alkoxycarbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄, NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl, R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,

009751766-122900

Sub B4
Cont.



wherein R_5' is selected from the group consisting of hydrogen, C_1-C_6 alkyl or aryl; R_{16}' is selected from hydrogen or one or two groups selected from C_1-C_6 alkyl, halogen, and C_1-C_6 alkoxy; Q is selected from the group consisting of -O-, -N(COR₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, CON(R₁₀), SO₂(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C_3-C_8 cycloalkyl or C_1-C_{10} alkyl; R₁₈ and R₁₉ are independently selected from the group consisting of hydrogen, unsubstituted C_1-C_{10} alkyl, substituted C_1-C_{10} alkyl, C_3-C_8 cycloalkyl, C_3-C_8 alkenyl, C_3-C_8 alkynyl and aryl or R₁₈ and R₁₉ may be combined with another element to which they are attached to form a radical Z having the formula



wherein Q₂ is selected from the group consisting of a covalent bond, -O-, -S-, -SO₂-, -CO-, -CO₂-, -N(C_1-C_6 alkyl)-, -N(CO C_1-C_6 alkyl)-, -N(SO₂ C_1-C_6 alkyl)-, -N(CO aryl)-, or -N(SO₂ aryl); R₂₀, R₂₁ and R₂₂ are independently selected from the group consisting of or C_1-C_6 alkyl; R₂₃ is selected from the group consisting of hydrogen, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, heteroaryl or aryl; wherein D is the residue of an active methylene compound selected from







09754766-122900

Sub B4
cont.


the group consisting of malononitrile, α -cyanoacetic acid esters, malonic acid esters, α -cyanacetic acid amides, α -C₁-C₆ alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles, α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles, α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane dinitrile-5,5-dioxides, 1,3-bis(dicyanomethylene) indanes, barbituric acid, 5-pyrazolones, dinedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides or aryl-C(CH₃)C=C(CN)₂, with the proviso that two acidic functional groups containing one acidic hydrogen each, or a functional group containing two acidic hydrogens are present.

58. The composition of claim 54 wherein the light absorbing portion of A₁ comprises the residue of at least one bis-azo light absorbing monomer wherein the bis coupling component Y₁ is represented by the structure Z₁-L₁-Z₂, wherein Z₁ and Z₂ are independently selected from the group consisting of

SUB
C17




or



- wherein, L_1 is bonded to the nitrogen atom of Z_1 and Z_2 ;
 wherein L_1 is selected from the group consisting of C_2 - C_{12}
 alkylene, C_3 - C_8 cycloalkylene, arylene, C_1 - C_4 alkylene-
- 5 C_3 - C_8 cycloalkylene- C_1 - C_4 alkylene, C_1 - C_4 alkylene-arylene-
 C_1 - C_4 alkylene, C_2 - C_4 alkylene-O-arylene-O- C_2 - C_4 alkylene,
 (C_2 - C_4 alkylene O)₁₋₃ C_2 - C_4 alkylene, C_2 - C_4 alkylene- S- C_2 - C_4
 alkylene, C_2 - C_4 alkylene-SO₂- C_2 - C_4 alkylene, C_2 - C_4
 alkylene-N(SO₂ C_1 - C_6 alkyl)- C_2 - C_4 alkylene, C_2 - C_4 alkylene-
- 10 N(SO₂ aryl)- C_2 - C_4 - alkylene, C_2 - C_4 alkylene- OCO₂- C_2 - C_4
 alkylene, C_2 - C_4 alkylene- O₂C-arylene-CO₂- C_2 - C_4 alkylene,
 C_2 - C_4 alkylene-O₂C- C_1 - C_{12} alkylene-CO₂- C_2 - C_4 alkylene, C_2 - C_4
 alkylene-O₂C- C_3 - C_8 cycloalkylene-CO₂- C_2 - C_4 alkylene, C_2 - C_4
 alkylene-NHCO- C_2 - C_4 alkylene and C_2 - C_4 alkylene-NHSO₂-
- 15 C_2 - C_4 alkylene; wherein R_{17} is selected from the group
 consisting of hydrogen, 1-2 groups selected from C_1 - C_6
 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, -O C_2 - C_6 alkylene-OH,
 O C_2 - C_6 alkylene- C_1 - C_6 alkanoyloxy, C_1 - C_6 alkylene-OH, C_1 - C_6
 alkylene- C_1 - C_6 alkanoyloxy, halogen, carboxy, C_1 - C_6
- 20 alkoxycarbonyl, trifluoromethyl, NHCOR₂₄ , NHCOR₂₄,
 NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R₂₄ is selected from the
 group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl
 or aryl, R₂₅ is selected from the group consisting of C_1 - C_{10}
 alkyl, C_3 - C_8 cycloalkyl or aryl wherein each C_1 - C_{10} alkyl
- 25 group in R₂₄ and R₂₅ may be further substituted with one or
 more groups selected from the group consisting of C_3 - C_8
 cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C_1 - C_6 alkyl,
 cyano, hydroxy, succinimido, C_1 - C_6 alkoxy,

09751766-122900

SUB
C17




5 wherein R_5' is selected from the group consisting of hydrogen, C_1 - C_6 alkyl or aryl; R_{16}' is selected from hydrogen or one or two groups selected from C_1 - C_6 alkyl, halogen and C_1 - C_6 alkoxy; Q is selected from the group consisting of -O-, -N(COR₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, CON(R₁₀), SO₂(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C_3 - C_8 cycloalkyl or C_1 - C_{10} alkyl; R₁₈ is selected from the group consisting of hydrogen, unsubstituted C_1 - C_{10} alkyl, substituted C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl and aryl; R₂₀, R₂₁ R₂₂ are independently selected from the group consisting of or C_1 - C_6 alkyl; R₂₃ is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, heteroaryl or aryl.


15 ~~59. The diacidic anthraquinone compounds having~~
Formulae

09751766-122900


SUB
C17




XIV




XIXc



XIXd




XIXe



XIXf

wherein R₁₄ is selected from the group consisting of
hydrogen, 1-4 groups selected from amino, C₁-C₁₀
alkylamino, C₃-C₈ alkenylamino, C₃-C₈ alkynylamino, C₃-C₈
5 cycloalkylamino, arylamino, halogen, C₁-C₆ alkoxy, C₁-C₆
alkylthio, aryl, aroyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy,
NHCO C₁-C₆ alkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHSO₂ C₁-C₆
alkyl, NHSO₂ aryl, C₁-C₆ alkoxy carbonyl, aryloxy, arylthio,
heteroarylthio, cyano, nitro, trifluoromethyl, thiocyno,
10 SO₂C₁-C₆ alkyl, SO₂ aryl, -SO₂NH C₁-C₆ alkyl, -SO₂N(C₁-C₆
alkyl)₂, -SO₂N(C₁-C₆ alkyl) aryl, CONH C₁-C₆ alkyl, CON(C₁-C₆
alkyl)₂, CON(C₁-C₆ alkyl) aryl, C₁-C₆ alkyl, furfurylamino,


tetrahydrofurfurylamino, 4-(hydroxymethyl)
cyclohexanemethylamino,



5

or hydroxy; Q and Q' are independently selected from the
group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-,
-S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is
10 selected from the group consisting of hydrogen, aryl, C₃-C₈
cycloalkyl, or C₁-C₁₀ alkyl; R₁₆' is selected from hydrogen
or one or two groups selected from C₁-C₆ alkyl, halogen and
C₁-C₆ alkoxy; wherein each C₁-C₆ alkyl group and C₁-C₆ alkyl
group which is a portion of another group may contain at
least one substituent selected from the group consisting
15 of hydroxy, cyano, chlorine, fluorine, C₁-C₆ alkoxy, C₃-C₈
cycloalkoxy, C₁-C₆ alkylcyclohexyl, hydroxymethyl
cyclohexyl, aryl and heteroaryl; with the provision that
two acidic groups containing one acidic proton each or one
acidic group containing two acidic hydrogens be present in
20 the compounds of Formula XIV, XIXc, XIXd, XIXe XIXf.

60. The diacidic anthraquinone compounds of claim
57 having the following structures:







25

09754766-122900

wherein Sub is a substituent selected from the group consisting of halogen, trifluoromethyl, aroyl, C₁-C₆ alkanoyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, aryloxy, arylthio, heteroarylthio, cyano, nitro, SO₂NHC₁-C₆ alkyl, SO₂N (C₁-C₆ alkyl)₂, SO₂N (C₁-C₆ alkyl) aryl, CONH C₁-C₆ alkyl, CON (C₁-C₆ alkyl)₂, CON (C₁-C₆ alkyl) aryl, C₁-C₆ alkyl, SO₂ C₁-C₆ alkylsulfonyl and SO₂ aryl; Sub₁ is a substituent selected from the group consisting of amino, C₁-C₁₂ alkylamino, arylamino and C₃-C₈ cycloalkylamino.

61. The diacidic anthrapyridone compounds having Formulae




wherein R₁₄ is selected from the group consisting of hydrogen, 1-4 groups selected from amino, C₁-C₁₀ alkylamino, C₃-C₈ alkenylamino, C₃-C₈ alkynylamino, C₃-C₈ cycloalkylamino, arylamino, halogen, C₁-C₆ alkoxy, C₁-C₆ alkylthio, aryl, aroyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy, NHCO C₁-C₆ alkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHSO₂ C₁-C₆ alkyl, NHSO₂ aryl, C₁-C₆ alkoxy, aryloxy, arylthio, heteroarylthio, cyano, nitro, trifluoromethyl, thiocyno, SO₂C₁-C₆ alkyl, SO₂ aryl, -SO₂NH C₁-C₆ alkyl, -SO₂N(C₁-C₆


09751766-122000

alkyl)₂, -SO₂N(C₁-C₆ alkyl) aryl, CONH C₁-C₆ alkyl, CON(C₁-C₆ alkyl)₂, CON(C₁-C₆ alkyl) aryl, C₁-C₆ alkyl or hydroxy; Q and Q' are independently selected from the group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-, -S-,
5 -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N (R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl, or C₁-C₁₀ alkyl; R₁₅ is selected from the group consisting of hydrogen, cyano, C₁-C₆ alkylamino, C₁-C₆ alkoxy, halogen, arylthio, aryl, heteroaryl,
10 heteroarylthio, C₁-C₆ alkoxycarbonyl, aroyl or arylsulfonyl; R₁₆ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl and aryl; R₁₆' is selected from the group consisting of hydrogen or one or two groups selected from C₁-C₆ alkyl, halogen and C₁-C₆ alkoxy; wherein each C₁-C₆ alkyl group and C₁-C₆ alkyl
15 group which is a portion of another group may contain at least one substituent selected from the group consisting of hydroxy, cyano, chlorine, fluorine, C₁-C₆ alkoxy, C₃-C₈ cycloalkoxy, C₁-C₆ alkylcyclohexyl, hydroxymethyl
20 cyclohexyl, aryl and heteroaryl; with the provision that two acidic groups containing one acidic proton each or one acidic group containing two acidic hydrogens be present in the compounds of Formula XVIII, XVIIIa, XVIIIb, and XIXa.


62. The diacidic anthrapyridone compound of claim
25 61 having the structure:



63. The diacidic anthraquinone compounds having the
30 formulae




or




5

where R_{16}' is selected from the group consisting of hydrogen or one or two groups selected from C_1-C_6 alkyl, halogen and C_1-C_6 alkoxy; and Sub_3 is a substituent selected from C_1-C_6 alkylthio, arylthio and heteroarylthio and Sub_2 is a substituent selected from the group consisting of amino, C_1-C_{10} alkylamino, C_3-C_8 alkenylamino, C_3-C_8 alkynylamino, C_3-C_8 cycloalkylamino, arylamino, furfurylamino, tetrahydrofurfurylamino, 4-(hydroxymethyl) cyclohexanemethylamino, $NHCO$ C_1-C_6 alkyl, $NHCO$ aryl, $NHCO_2$ C_1-C_6 alkyl, $NHSO_2$ C_1-C_6 alkyl, $NHSO_2$ aryl and




20

64. The diacidic anthraquinone compounds of claim 59 having the formulae:




wherein Sub₂ is as defined in claim 63; Sub₄ is selected from the group consisting of Sub₂, NHCO C₁-C₆ alkyl, NHCO₂ C₁-C₆ alkyl, NHCO aryl, NHSO₂ C₁-C₆ alkyl, NHSO₂ aryl, C₁-C₆ alkylthio, arylthio, heteroarylthio and hydroxy; Q is selected from the group consisting of -O-, S-, -SO₂-; Q' selected from the group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl, or C₁-C₁₀ alkyl.

65. A diacidic anthraquinone compounds having the formula




wherein Sub, Sub₁ and R₁₆' are as defined in claim 60.

66. The diacidic anthraquinone compounds having the structures




- wherein Q is selected from the group consisting of -O-, -S- and -SO₂-; Q' is selected from the group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl, or C₁-C₁₀ alkyl; and R₁₆' is selected from the group consisting of hydrogen or one or two groups selected from C₁-C₆ alkyl, halogen and C₁-C₆ alkoxy.

67. The diacidic anthraquinone compounds having the structures:



wherein Sub₁ defined as in claim 60, Sub₄ is defined as in claim 64, Q is selected from the group consisting of -O-,
 5 -S- and -SO₂-; Q' is selected from the group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl, or C₁-C₁₀ alkyl; and R₁₆' is selected from the group consisting of hydrogen or one or two groups selected from C₁-C₆ alkyl, halogen and C₁-C₆ alkoxy.

68. The diacidic anthraquinone compounds having the structures:



wherein Q is selected from the group consisting of -O-, -S- and -SO₂-; Sub₁ is a substituent selected from the group consisting of amino, C₁-C₁₂ alkylamino, arylamino and C₃-C₈ cycloalkylamino; Q' is selected from the group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl, or C₁-C₁₀ alkyl; and R₁₆' is selected from the group consisting of hydrogen or one or two groups selected from C₁-C₆ alkyl, halogen and C₁-C₆ alkoxy.

69. The diacidic azo and disazo compounds of the formulae R₆-N=N-Z (VI) and R₆-N=N-R₇-N=N-Z (VII), respectively wherein R₆ is the residue of a diazotized aromatic or heteroaromatic amine and Z is the residue of an electron rich coupling component selected from the group consisting of the classes of anilines, 1-aminonaphthalenes, 1,2-dihydroquinolines, 1,2,3,4-tetrahydroquinolines, benzomorpholines (3,4-dihydro-2H-1,4-benzoxazine), pyrazolones, pyrazoles, 3-cyano-6-hydroxy-2-pyridones, 2,3-dihydroindoles, indoles, 4-hydroxycoumarins, 4-hydroxy-2-quinolones, imidazo[2,1-b]thiazoles, julolidines (2,3,6,7-tetrahydro-1H,5H-benzo[ij]quinolizines), 1-oxajulolidines, 1,2,5,6-tetrahydro-4H-pyrrolo[3,2,1-ij]quinolines, 2,6-diamino-3-cyanopyridines, 2-aminothiazoles, 2-aminothiophenes, 5,5-dimethyl-1,3-cyclohexanedione (dimedone), phenols, naphthols, 2,4-pentanediones or acetoacetarilides; R₇ is a divalent aromatic or heteroaromatic radical selected from the group consisting of the classes of 1,4-phenylene, naphthalene -1, 4-diyl, thiazol-2,5-diyl and thiophene -

00622T 9975760

2,5-diyl; with the provision that R₆ or Z contains a carboxy (-CO₂H) acidic group and that another acidic group selected from the group consisting of (-CO₂H) , -SH, -OH attached to aromatic ring, -CONHCO-, -SO₂NH-CO-, -SO₂NH-SO₂-, and 1(H) 1, 2, 4-triazol-3-yl, be present on or as part of R₆ or Z so that each R₆ and Z moiety contains one acidic group.

70. The diacidic azo and disazo compounds of claim 69 wherein R₆ and Z each contain a carboxy (-CO₂H) acidic group.


71. The diacidic azo and bisazo compounds of claim 69 wherein R₆ is the residue of substituted diazotized aromatic or heteroaromatic amine compounds derived from the classes of aniline, 1-aminonaphthalene, 1-aminoanthraquinone, 4-aminoazobenzene, 2-aminothiazole, 2-aminobenzothiazole, 3-amino- 2,1-benzisothiazole, 2-aminothieno[2,3-d]thiazole, 5-aminoisothiazole, 5-aminopyrazole, 4-aminopyrazoloisothiazole, 2-amino-1,3,4-thiadiazole, 5-amino-1,2,4-thiadiazole, 5-amino-1,2,3-triazole, 2-amino-1,3,4-triazole, 2(5) aminoimidazole, 3-aminopyridine, 2(3) aminothiophene, 2(3)-aminobenzo[b]thiophene, 2-aminothieno[3,2-b]thiophene, 3-aminothieno[2,3-c]isothiazole, 3-amino-7-benz-2,1-isothiazole, 3-aminobenzothienoisothiazole, 3-aminoisothiazole[3,4-d]pyrimidine, 5-amino-1,2,3-triazole, 3(4) aminophthalimide and 5(6) amino-1,2-benzisothiazolon-1,1-dioxide and Z is the residue of an electron rich coupling coupler residue selected from the group consisting of the following:

0951766-13300




wherein R₁₇ is selected from the group consisting of hydrogen, 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆

- alkoxy, C₁-C₆ alkylthio, -O C₂-C₆ alkylene-OH, O C₂-C₆ alkylene- C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene- C₁-C₆ alkanoyloxy, halogen, carboxy, C₁-C₆ alkoxy, carbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄,
 5 NHCON(R₂₄)R₂₅, and NHCO₂R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl, R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or
 10 more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,



- 15 wherein R₅', R₁₆' and Q are as defined in claim 63; R₁₈ and R₁₉ are independently selected from the group consisting of hydrogen, unsubstituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl and
 20 aryl or R₁₈ and R₁₉ may be combined with another element to which they are attached to form a radical Z having the formula



wherein Q_2 is selected from the group consisting of a covalent bond, -O-, -S-, -SO₂-, -CO-, -CO₂-, -N-(C₁-C₆ alkyl)-, -N(CO C₁-C₆ alkyl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(CO aryl)-, or -N(SO₂ aryl); R_{20} , R_{21} and R_{22} are independently
5 selected from the group consisting of or C₁-C₆ alkyl; R_{23} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, heteroaryl or aryl.

72. The diacidic azo and disazo compounds of claim 69 of the formulae $R_6-N=N-Z$ (VI) and $R_6-N=N-R_7-N=N-Z$ (VII),
10 respectively, wherein one of R_6 and Z contains two carboxy (-CO₂H) acidic groups.


73. The diacidic azo and disazo compounds of claim 69 or 72 wherein R_6 is the residue of a diazotized substituted or unsubstituted diazotized aromatic or
15 heteromatic amine compound derived from an amine selected from aniline, 1-aminonaphthalene, 1-aminoanthraquinone, 4-aminoazobenzene, 2-aminothiazole, 2-aminobenzothiazole, 3-amino- 2,1-benzisothiazole, 2-aminothieno[2,3-d]thiazole, 5-aminoisothiazole, 5-aminopyrazole,
20 4-aminopyrazoloisothiazole, 2-amino-1,3,4-thiadiazole, 5-amino-1,2,4-thiadiazole, 5-amino-1,2,3-triazole, 2-amino-1,3,4-triazole, 2(5) aminoimidazole, 3-aminopyridine, 2(3) aminothiophene, 2(3) aminobenzo[b]thiophene, 2-aminothieno[3,2-b]thiophene, 3-aminothieno[2,3-
25 c]isothiazole, 3-amino-7-benz- 2,1-isothiazole, 3-aminobenzothienoisothiazole, 3-aminoisothiazole[3,4-d]pyrimidine, 5-amino- 1,2,3-triazole, 3(4) aminophthalimide and 5(6) amino-1,2-benzisothiazolon-1,1-dioxide and Z is the residue of an electron rich coupling
30 component selected from the group consisting of the following:

09751766-122900




wherein R₁₇ is selected from the group consisting of hydrogen, 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆

- alkoxy, C₁-C₆ alkylthio, -O C₂-C₆ alkylene-OH, O C₂-C₆ alkylene- C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene- C₁-C₆ alkanoyloxy, halogen, carboxy, C₁-C₆ alkoxycarbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄,
 5 NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl, R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or
 10 more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,




- 15 wherein R₅', R₁₆' and Q are as defined in claim 63; R₁₈ and R₁₉ are independently selected from the group consisting of hydrogen, unsubstituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl and
 20 aryl or R₁₈ and R₁₉ may be combined with another element to which they are attached to form a radical Z having the formula




wherein Q_2 is selected from the group consisting of a covalent bond, -O-, -S-, -SO₂-, -CO-, -CO₂-, -N-(C₁-C₆ alkyl)-, -N(CO C₁-C₆ alkyl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(CO aryl)-, or -N(SO₂ aryl); R_{20} , R_{21} and R_{22} are independently selected from the group consisting of or C₁-C₆ alkyl; R_{23} is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₁-C₈ cycloalkyl, heteroaryl or aryl.

74. The diacidic azo and bisazo compounds of claim 72 wherein Z is an electron rich coupler selected from the group consisting of the following:




006227-9975760




- wherein R_{17} is selected from the group consisting of
- 5 hydrogen, 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, $-O$ C_2 - C_6 alkylene-OH, O C_2 - C_6 alkylene- C_1 - C_6 alkanoyloxy, C_1 - C_6 alkylene-OH, C_1 - C_6 alkylene- C_1 - C_6 alkanoyloxy, halogen, carboxy, C_1 - C_6 alkoxy, carbonyl, trifluoromethyl, $NHCO_2R_{24}$, $NHCO_2R_{24}$,
 - 10 $NHCON(R_{24})R_{25}$, and $NHSO_2R_{25}$, wherein R_{24} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl or aryl, R_{25} is selected from the group consisting of C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl or aryl wherein each C_1 - C_{10} alkyl group in R_{24} and R_{25} may be further substituted with one or
 - 15 more groups selected from the group consisting of C_3 - C_8 cycloalkyl, aryl, aryloxy, arylthio, CO_2H , CO_2 C_1 - C_6 alkyl, cyano, hydroxy, succinimido, C_1 - C_6 alkoxy,

006221 9975166 122900



wherein R_5' , R_{16}' and Q are as defined in claim 63; R_{18} and R_{19} are independently selected from the group consisting of hydrogen, unsubstituted C_1 - C_{10} alkyl, substituted C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl and aryl or R_{18} and R_{19} may be combined with another element to which they are attached to form a radical Z having the formula



wherein Q_2 is selected from the group consisting of a covalent bond, -O-, -S-, -SO₂-, -CO-, -CO₂-, -N-(C_1 - C_6 alkyl)-, -N(CO C_1 - C_6 alkyl)-, -N(SO₂ C_1 - C_6 alkyl)-, -N(CO aryl)-, or -N(SO₂ aryl); R_{20} , R_{21} and R_{22} are independently selected from the group consisting of or C_1 - C_6 alkyl; R_{23} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, heteroaryl or aryl.


75. The diacidic bisazo compounds having the formula R_6 -N=N-Y₁-N=N-R₆ (VIIa) wherein R_6 is the residue of a substituted diazotized aromatic or heteroaromatic amine compound derived from the classes of aniline, 1-aminonaphthalene, 1-aminoanthraquinone, 4-aminoazobenzene, 2-aminothiazole, 2-aminobenzothiazole, 3-amino-2,1-benzisothiazole, 2-aminothieno[2,3-d]thiazole,

09751766-132900

5-aminoisothiazole, 5-aminopyrazole,
4-aminopyrazoloisothiazole, 2-amino-1,3,4-thiadiazole, 5-
amino-1,2,4-thiadiazole, 5-amino-1,2,3-triazole, 2-amino-
1,3,4-triazole, 2(5) aminoimidazole, 3-aminopyridine, 2(3)
5 aminothiophene, 2(3) aminobenzo[b]thiophene, 2-
aminothieno[3,2-b]thiophene, 3-aminothieno[2,3-
c]isothiazole, 3-amino-7-benz- 2,1-isothiazole, 3-
aminobenzothienoisothiazole, 3-aminoisothiazole[3,4-
d]pyrimidine, 5-amino- 1,2,3-triazole, 3(4)
10 aminophthalimide and 5(6) amino-1,2-benzisothiazolon-1,1-
dioxide and Y_1 is the residue of a bis coupling component
selected from the group consisting of the classes of
anilines, 1,2-dihydroquinolines, 1,2,3,4-
tetrahydroquinolines, benzomorpholines (3,4-dihydro-
15 2H-1,4-benzoxazines), 3-cyano-6-hydroxy-2-pyridones, 2,6-
diaminopyridines, 2,3-dihydroindoles, naphthylamines, 2-
aminothiazoles, or a combination of these, with the
provision that each R_6 group contain one acidic group
selected from the group consisting of $-CO_2H$, $-SH$, $-OH$
20 attached to an aromatic ring, $-NHCONH-$, $-SO_2NHCO-$, $-$
 SO_2NHSO_2- , 1 (H)-1,2,4-triazol-3-yl-, imidazolyl,
benzimidazolyl, pyrazolyl and $-SO_2H$ attached to aromatic
ring.

76. The bis-azo compounds of claim 75 wherein each
25 R_6 group contains one carboxy ($-CO_2H$) group.


77. The bis-azo compounds of claim 75 wherein Y_1 has
the formula $Z_1-L_1-Z_2$ wherein Z_1 and Z_2 are independently
selected from the group consisting of:



wherein L_1 is bonded to the nitrogen atom of Z_1 and Z_2 ;
 wherein L_1 is selected from the group consisting of C_2 - C_{12}
 5 alkylene, C_3 - C_8 cycloalkylene, arylene, C_1 - C_4 alkylene-
 C_3 - C_8 cycloalkylene- C_1 - C_4 alkylene, C_1 - C_4 alkylene-arylene-
 C_1 - C_4 alkylene, C_2 - C_4 alkylene-O-arylene-O- C_2 - C_4 alkylene,
 (C_2 - C_4 alkylene O)₁₋₃ C_2 - C_4 alkylene, C_2 - C_4 alkylene- S- C_2 - C_4
 alkylene, C_2 - C_4 alkylene-SO₂- C_2 - C_4 alkylene, C_2 - C_4
 10 alkylene-N(SO₂ C_1 - C_6 alkyl)- C_2 - C_4 alkylene, C_2 - C_4 alkylene-
 N(SO₂ aryl)- C_2 - C_4 - alkylene, C_2 - C_4 alkylene-OCO₂- C_2 - C_4
 alkylene, C_2 - C_4 alkylene- O₂C-arylene-CO₂- C_2 - C_4 alkylene,
 C_2 - C_4 alkylene-O₂C- C_1 - C_{12} alkylene-CO₂- C_2 - C_4 alkylene, C_2 - C_4
 alkylene-O₂C- C_3 - C_8 cycloalkylene-CO₂- C_2 - C_4 alkylene, C_2 - C_4
 15 alkylene-NHCO- C_2 - C_4 alkylene and C_2 - C_4 alkylene-NHSO₂-
 C_2 - C_4 alkylene; R_{17} is selected from the group consisting
 of hydrogen, 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6
 alkoxy, C_1 - C_6 alkylthio, -O C_2 - C_6 alkylene-OH, O C_2 - C_6
 alkylene- C_1 - C_6 alkanoyloxy, C_1 - C_6 alkylene-OH, C_1 - C_6
 20 alkylene- C_1 - C_6 alkanoyloxy, halogen, carboxy, C_1 - C_6
 alkoxy, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄,
 NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R_{24} is selected from the
 group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl
 or aryl, R_{25} is selected from the group consisting of C_1 - C_{12}

005221" 9975460








alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,



wherein R₅', R₁₆' and Q are as defined in claim 63; R₁₈ is selected from the group consisting of hydrogen, a group selected from the group consisting of unsubstituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl and aryl; R₂₀, R₂₁ and R₂₂ are independently selected from the group consisting of or C₁-C₆ alkyl; R₂₃ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, heteroaryl or aryl.

78. The diacidic methine, arylidene, polymethine, azamethine, 3-aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene -2-oxypyrroline and aryl isoindoline corresponding to formulae VIII, VIIIA, VIIIB, IX, X, XI and XII, respectively:

006227 997560




wherein R_{11} is the residue of an aniline, 1-naphthylamine, 1,2-dihydroquinoline, 1,2,3,4-tetrahydroquinoline, 1,3,3-trimethyl-2-methyleneindole, 1,3-dihydro-2-methylene-1,1,3-trimethyl-2H-benz[e]indole, imidazo [2,1-b]thiazole, benzomorpholine (3,4-dihydro-2H-1,4-benzoxazine), indole, 2,3-dihydroindole, 2-aminothiazole, julolidine (2,3,6,7-tetrahydro-1H, 5H- benz [ij] quinolizine, 1-oxajulolidine, 4H-pyrrolo [3,2,1-ij]-quinoline, phenol, naphthol, thiophenol, pyrrole, pyrazole, furan, thiophene, carbazole, phenothiazine or phenoxazine compound; R_{12} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 alkenyl, C_3 - C_8 -alkynyl, C_3 - C_8 cycloalkyl, aryl, $(\text{CH}_2\text{CH}_2\text{O})_{1-3}$ R_{13} and C_1 - C_4 alkylene- C_3 - C_8 cycloalkylene, wherein the C_1 - C_6 alkyl groups may be substituted by at least one group selected from the group consisting of carboxy, C_1 - C_6 carbalkoxy, C_1 - C_6 alkanoyloxy, cyano, hydroxy, chlorine, fluorine, C_1 - C_6 alkoxy, C_3 - C_8 cycloalkyl or aryl; R_{13} is selected from the group consisting of hydrogen, C_1 - C_6 alkoxy or C_1 - C_6 alkanoyloxy; wherein D is the residue of an active

09751766-122900


- methylene compound selected from the group consisting of malononitrile, α -cyanoacetic acid esters, malonic acid esters, α -cyanacetic acid amides, α -C₁-C₆ alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles, 5 α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles, α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane 10 dinitrile-5,5-dioxides, 1,3-bis (dicyanomethylene) indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides or aryl-C(CH₃)C=C(CN)₂, with the proviso that two acidic functional groups selected from the group consisting of -CO₂H, -SH, - 15 OH attached to aromatic ring, -CONHCO-, -SO₂NHCO-, -SO₂NHSO₂-, 1(H) -1,2,4,-triazol-3-yl, imidazolyl, benzimidazolyl, pyrazolyl and SO₂H attached to aromatic ring be present or one diacidic sulfamoyl (-SO₂NH₂) be present.
- 20 79. The diacidic anthrapyridine compounds having the structures:

09751766-132900



wherein Sub₅ is a substituent selected from the group
 5 consisting of -N (C₁-C₁₀ alkyl)₂, -N (C₁-C₁₀ alkyl) aryl, -
 N (C₁-C₁₀ alkyl) C₃-C₈ cycloalkyl, morpholino and
 piperidino; Q and Q' are selected from the group
 consisting of -NH-, -O-, -S- and -SO₂-, R₁₆' is selected
 10 from the group consisting of hydrogen, C₁-C₆ alkyl, halogen
 and C₁-C₆ alkoxy.

80. The nitroarylamine compounds having the
 structure:



15 wherein ring A may be substituted with one or more groups
 selected from the group consisting of halogen, -SO₂N(C₁-C₆
 alkyl)₂, -CON (C₁-C₆ alkyl)₂, SO₂ C₁-C₆ alkyl, SO₂ aryl, C₁-
 C₆ alkyl, C₁-C₆ alkoxy, carboxy, and nitro; Ar is
 phthalimid-3 (or 4)-yl, phenyl, or 2-thienyl, or these
 20 substituted with one or more groups selected from the


006237 99475260

group consisting of halogen, C₁-C₆ alkyl, C₁-C₆ alkoxy, 1(H)-1,2,4,-triazolyl-3-ylthio, carboxy, or hydroxy, with the provision that two acidic groups be present.

81. The nitroarylamine compounds of claim 80 wherein two carboxy groups are present on Ar or ring A or one carboxy is present on each of Ar and ring A.

82. The diacidic compounds of claim 72, having the formula R₆-N=N-Z, wherein R₆ is selected from the group consisting of

10




wherein Sub₆ is selected from the group consisting of hydrogen, one to four groups selected from C₁-C₁₀ alkyl, C₁-C₆ alkoxy, C₃-C₈ cycloalkyl, halogen, , C₁-C₆ alkoxycarbonyl, formyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy, dicyanovinyl, C₃- C₈-cycloalkanoyl, thiocyno, trifluoroacetyl, cyano, carbamoyl, -CONH C₁-C₆ alkyl, CONHaryl, CON(C₁-C₆ alkyl)₂, SO₂N(C₁-C₆ alkyl)₂, CONH C₃-C₈ cycloalkyl, aryl, aroyl, -N(C₁-C₆ alkyl)SO₂ C₁-C₆ alkyl, NHCO C₁-C₆ alkyl, NHCO C₃-C₈ cycloalkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHCONH C₁-C₆ alkyl, NHCONHaryl, N(C₁-C₆ alkyl)aryl, arylazo, heteroaryl, aryloxy, arylthio, C₃-C₈ cycloalkoxy, heteroarylazo, heteroarylthio, arylsulfonyl, tricyanovinyl, aryloxysulfonyl, C₁-C₆ alkylsulfonyl, trifluoromethyl, fluorosulfonyl, trifluoromethylsulfonyl, thiocyno, nitro and CH=D, wherein D is the residue of an active methylene compound selected from the group consisting of malononitrile, α-cyanoacetic acid esters, malonic acid esters, α-cyanacetic acid amides, α-C₁-C₆ alkylsulfonylacetonitriles, α-arylsulfonylacetonitriles, α-C₁-C₆ alkanoylacetonitriles, α-aroylacetonitriles, α-heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-

09751766 1.22900



indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane dinitrile-5,5-dioxides, 1,3-bis (dicyanomethylene)

- 5 indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides and aryl- $C(CH_3)C(CN)_2$; with the provision that one acidic group selected from the group consisting of carboxy,




be present on either R_{17} , R_{18} , R_{19} or R_{24} ; Q is selected from the group consisting of -O-, -S-, and -SO₂-.


83. The compounds of claim 82 wherein R_{18} is
15 selected from the group consisting of C₁-C₁₀ alkylene-CO₂H,




84. The diacidic compounds of claim 82 wherein R_6 is
20 selected from the group consisting of




and Z is selected from



and




wherein R_{17} is selected from the group consisting of
 hydrogen, 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6
 5 alkoxy, halogen, $NHCO_2R_{24}$, $NHCONHR_{24}$, wherein
 R_{24} is selected from the group consisting of hydrogen, C_1 - C_{10}
 C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl and aryl; wherein each C_1 - C_{10}
 alkyl group in R_{24} may be further substituted with one or
 more groups selected from the group consisting of C_3 - C_8
 10 cycloalkyl, carboxy, aryl, aryloxy, arylthio, CO_2 C_1 - C_6
 succinimido, C_1 - C_6 alkoxy and
 alkyl, cyano, hydroxy,




are independently selected from
 the group consisting of hydrogen, unsubstituted C_1 - C_{10}
 alkyl, substituted C_1 - C_{10} alkyl, C_3 - C_8 alkyl, C_3 - C_8 alkenyl
 15 and aryl or R_{18} and R_{19} may be combined with another
 element to which they are attached to from a radical

00954756 " 12900





wherein Q_2 is selected from the group consisting of -O-, -S-, -SO₂-, -CO-, -CO₂-, -N (COC₁-C₆ alkyl)-, -N (SO₂ C₁-C₆ alkyl)-, -N (COaryl)-, and -N (SO₂ aryl)-; R_{20} , R_{21} , and R_{22} are independently selected from the group consisting of or C₁-C₆ alkyl; with the provision that either R_{17} contain one acidic group selected from the group consisting of carboxy and

10



with the groups R_{18} and R_{19} being void of acidic groups or R_{17} may be void of acidic groups and R_{18} be selected from the group consisting of C₁-C₁₀ alkylene -CO₂H,

15




wherein Q is selected from the group consisting of -O-, -S-, and -SO₂-; with the final provision that only two carboxy groups be present.

20

85. The diacidic compounds of claim 61 having the formula $R_6-N=N-Z$, wherein R_6 is the residue of a

09751766-122900

[illegible]

substituted heterocyclic diazotized amine and selected from the group consisting of

The diagram shows several chemical structures of substituted heterocyclic diazotized amines. These include:


- A thiazole ring with a carboxylic acid group (CO_2H) and a substituent Sub_7 .
- An imidazole ring with a carboxylic acid group (CO_2H) and a substituent Sub_8 .
- A pyrazole ring with two carboxylic acid groups (HO_2C) and a substituent Sub_9 .
- A thiadiazole ring with a carboxylic acid group (HO_2C) and a substituent Sub_{10} .
- A benzothiazole derivative with a carboxylic acid group (HO_2C) and a substituent Sub_{11} .
- A pyridine-like structure with a carboxylic acid group (HO_2C) and a substituent Q .


wherein Sub_7 is one or more substituent selected from the group consisting of hydrogen, halogen, $\text{C}_1\text{-C}_6$ alkyl, heteroaryl, $\text{C}_1\text{-C}_6$ alkanoyl, $\text{CONH C}_1\text{-C}_6$ alkyl, $\text{SO}_2 \text{C}_1\text{-C}_6$ alkyl, SO_2 aryl, $\text{C}_1\text{-C}_6$ alkoxy carbonyl, aroyl, cyano, and nitro; Sub_8 is selected from the group consisting of hydrogen, halogen, $\text{C}_1\text{-C}_6$ alkyl, aryl and heteroaryl; Sub_9 is selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, cycloalkyl, $\text{C}_3\text{-C}_8$ alkenyl, and aryl; Sub_{10} is selected from the group consisting of cyano, nitro, $\text{C}_1\text{-C}_6$ alkyl, arylthio, arylsulfonyl and $\text{C}_1\text{-C}_6$ alkoxy carbonyl; Sub_{11} is selected from the group consisting of hydrogen one or more substituents selected from the group consisting of $\text{C}_1\text{-C}_6$ alkyl, cyano, nitro, halogen,

heteroarylthio, arylsulfonyl, aryloxy and C₁-alkylsulfonyl; Q is selected from the group consisting of -O-, -S- and -SO₂-; wherein Z is a coupling component selected from the group consisting of


The image shows several chemical structures, many of which are crossed out with a large diagonal line. The structures include:

- A benzene ring with substituents R₁₇ and R₁₈ on the nitrogen atom.
- A quinoline derivative with substituents R₁₇, R₁₈, R₂₀, R₂₁, and R₂₂.
- A benzimidazole derivative with substituents R₁₇, R₁₈, R₂₀, and R₂₃.
- A benzoxazine derivative with substituents R₁₇, R₁₈, R₂₀, and R₂₁.
- A benzimidazole derivative with substituents R₁₇, R₁₈, R₂₀, and R₂₁.
- A benzimidazole derivative with substituents R₁₇, R₁₈, R₂₀, and R₂₁.







- 5 wherein R_{17} is selected from the group consisting of hydrogen, 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, C_1 - C_6 alkylthio, $-O$ C_2 - C_6 alkylene-OH, O C_2 - C_6 alkylene- C_1 - C_6 alkanoyloxy, C_1 - C_6 alkylene-OH, C_1 - C_6 alkylene- C_1 - C_6 alkanoyloxy, halogen, C_1 - C_6 alkoxy, carbonyl, trifluoromethyl, $NHCO_2R_{24}$, $NHCO_2R_{24}$, and $NHCONHR_{24}$, wherein
- 10 R_{24} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl and aryl; wherein each C_1 - C_{10} alkyl group in R_{24} may be further substituted with one or more groups selected from the group consisting of C_3 - C_8
- 15 cycloalkyl, aryl, aryloxy, arylthio, CO_2 C_1 - C_6 alkyl, cyano, hydroxy, succinimido, C_1 - C_6 alkoxy,



wherein R_5' is selected from the group consisting of hydrogen, C_1-C_6 alkyl and aryl; R_{16}' is selected from the group consisting of hydrogen, one or two groups selected from C_1-C_6 alkyl, halogen and C_1-C_6 alkoxy; R_{18} and R_{19} are independently selected from the group consisting of hydrogen, unsubstituted C_1-C_{10} alkyl, substituted C_1-C_{10} alkyl, C_3-C_8 cycloalkyl, C_3-C_8 alkenyl, C_3-C_8 alkynyl and aryl or R_{18} and R_{19} may be combined with another element to which they are attached to form a radical Z having the formula





wherein Q_2 is selected from the group consisting of a covalent bond, $-O-$, $-S-$, $-SO_2-$, $-CO-$, $-CO_2-$, $-N-(C_1-C_6 \text{ alkyl})-$, $-N(CO \ C_1-C_6 \text{ alkyl})-$, $-N(SO_2 \ C_1-C_6 \text{ alkyl})-$, $-N(CO \ aryl)-$, or $-N(SO_2 \ aryl)-$; R_{20} , R_{21} and R_{22} are independently selected from the group consisting of or C_1-C_6 alkyl; R_{23} is selected from the group consisting of hydrogen, C_1-C_6 alkyl, C_3-C_8 cycloalkyl, heteroaryl or aryl; with the provision that one acidic group selected from the group consisting of carboxy,




be present on either R_{17} , R_{18} , R_{19} or R_{24} ; Q is selected from the group consisting of $-O-$, $-S-$, and $-SO_2-$; with the provision that R_6 and Z each contain one acidic group.


86. The diacidic compounds of claim 85 having the formula $R_6-N=N-Z$, wherein R_6 is selected from the group consisting of




and Z is selected from the group consisting of




wherein R_{17} is selected from the group consisting of hydrogen, 1-2 groups selected from C_1 - C_6 alkyl, C_1 - C_6 alkoxy, halogen, $NHCO_2R_{24}$, $NHCONHR_{24}$, wherein R_{24} is selected from the group consisting of hydrogen, C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl and aryl; wherein each C_1 - C_{10} alkyl group in R_{24} may be further substituted with one or more groups selected from the group consisting of C_3 - C_8 cycloalkyl, carboxy, aryl, aryloxy, arylthio, CO_2 C_1 - C_6 alkyl, cyano, hydroxy, succinimido, C_1 - C_6 alkoxy and



are independently selected from the group consisting of hydrogen, unsubstituted $\text{C}_1\text{-C}_{10}$ alkyl, substituted $\text{C}_1\text{-C}_{10}$ alkyl, $\text{C}_3\text{-C}_8$ alkyl, $\text{C}_3\text{-C}_8$ alkenyl and aryl or R_{18} and R_{19} may be combined with another
5 element to which they are attached to from a radical





wherein Q_2 is selected from the group consisting of $-\text{O}-$, $-\text{S}-$, $-\text{SO}_2$, $-\text{CO}-$, $-\text{CO}_2$, $-\text{N}(\text{CO}\text{C}_1\text{-C}_6\text{ alkyl})-$, $-\text{N}(\text{SO}_2\text{ C}_1\text{-C}_6\text{ alkyl})-$, $-\text{N}(\text{COaryl})-$, and $-\text{N}(\text{SO}_2\text{ aryl})-$; R_{20} , R_{21} , and R_{22} are independently selected from the group consisting of or $\text{C}_1\text{-C}_6$ alkyl; with the provision that either R_{17} contain one
10 acidic group selected from the group consisting of carboxy
15 and




with the groups R_{18} and R_{19} being void of acidic groups or
20 R_{17} may be void of acidic groups and R_{18} be selected from the group consisting of $\text{C}_1\text{-C}_{10}$ alkylene $-\text{CO}_2\text{H}$,

006221 997560




wherein Q is selected from the group consisting of - O -,
- S -, and - SO₂ -; with the final provision that only two
5 carboxy groups be present.


87. The diacidic compounds of claim 71 having the
formula R₆-N=N-Z, wherein R₆ is the residue of a diazotized
aromatic amine and is selected from the group consisting
of



wherein Sub₁₂ is one or more groups selected from the group
consisting of hydrogen, C₁-C₆ alkyl, cyano, nitro, C₁-C₆
15 alkylthio, C₁-C₆ alkylsulfonyl, aryl, heteroaryl, arylthio,
arylsulfonyl, halogen, trifluoromethyl, alkanoyl, aroyl,
formyl, NHCO aryl, NHCO C₁-C₆ alkyl, C₁-C₆ alkoxy, carbonyl,
C₁-C₆ alkoxy and - SO₂N (C₁-C₆ alkyl)₂; Q is selected from
the group consisting of -O-, -S-, -SO₂-, -CONH- and -SO₂N
20 (C₁-C₆ alkyl)-; Z is selected from the group consisting of




alkoxy, C₁-C₆ alkylthio, -O- C₂-C₆ alkylene - OH, O-C₂-C₆ alkylene- C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene - C₁-C₆ alkanoyloxy, halogen, C₁-C₆ alkoxycarbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄ and NHCON(R₂₄) R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl and aryl; R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl and aryl; wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido and C₁-C₆ alkoxy; R₁₈ and R₁₉ are selected from the group consisting of hydrogen C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl, and aryl; R₁₈ and R₁₉ in combination may be combined with another element to which they are attached to form a radical Z having the formula




wherein Q₂ is selected from the group consisting of a covalent bond, -O-, -SO₂-, -S-, -CO-, -CO₂-, -N(COC₁-C₆ alkyl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(CO aryl)-, -N(SO₂ aryl)-; R₂₀, R₂₁ and R₂₂ are independently selected from the group consisting of or C₁-C₆ alkyl; R₂₃ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl, heteroaryl or aryl; with the provision that no acidic groups be present on Z.


88. The diacidic compounds of claim 87 wherein R₆ has the structure



and Z is selected from




89. The diacidic compound of claim 72 wherein R₆ is residue of a diazotized heterocyclic amine and is selected from the group consisting of



wherein Sub₈ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, halogen, aryl and heteroaryl; Sub₉ is selected from the group consisting of C₁-C₆ alkyl C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, and aryl; Q is selected from the group consisting of -O-, -S- and -SO₂-; with the provision that no acidic groups be present on Z.

90. The diacidic compounds of claim 72 wherein R₆ is selected from the group consisting of



5

wherein Sub₈ is selected from the group consisting of hydrogen, C₁-C₆ alkyl, heteroaryl and aryl; Sub₉ is selected from the group consisting of C₁-C₆ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl and aryl; Sub₁₃ is selected from the group consisting of hydrogen, one or more groups selected from C₁-C₁₀ alkyl, C₁-C₆ alkoxy, C₃-C₈ cycloalkyl, halogen, , C₁-C₆ alkoxycarbonyl, formyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy, dicyanovinyl, C₃- C₈-cycloalkanoyl, thiocyno, trifluoroacetyl, cyano, carbamoyl, -CONH C₁-C₆ alkyl, CONHaryl, CON(C₁-C₆ alkyl)₂, SO₂N(C₁-C₆ alkyl)₂, CONH C₃-C₈ cycloalkyl, aryl, aroyl, -N(C₁-C₆ alkyl)SO₂ C₁-C₆ alkyl, NHCO C₁-C₆ alkyl, NHCO C₃-C₈ cycloalkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHCONH C₁-C₆ alkyl, NHCONHaryl, N(C₁-C₆

10


15


09751766-122900

alkyl)aryl, arylazo, heteroaryl, aryloxy, arylthio, C₃-C₈
cycloalkoxy, heteroarylazo, heteroarylthio, arylsulfonyl,
tricyanovinyl, aryloxysulfonyl, C₁-C₆ alkylsulfonyl,
trifluoromethyl, fluorosulfonyl, trifluoromethylsulfonyl,
5 thiocyano, nitro and CH=D, wherein D is the residue of an
active methylene compound selected from the group
consisting of malononitrile, α-cyanoacetic acid esters,
malonic acid esters, α-cyanoacetic acid amides, α-C₁-C₆
alkylsulfonylacetonitriles, α-arylsulfonylacetonitriles,
10 α-C₁-C₆ alkanoylacetonitriles, α-arylacetonitriles,
α-heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-
indanediones, 2-furanones, benzo-2-furanones, naphtho-2-
furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-
dioxo (2H)-pyridines, benzo (b) thieno-3-ylidene propane
15 dinitrile-5,5-dioxides, 1,3-bis (dicyanomethylene)
indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-
2,3-dihydro-1-benzothiophene-1,1-dioxides and aryl-
C(CH₃)C=C(CN)₂; Sub₁₄ is selected from the group consisting
of hydrogen, halogen, cyano, C₁-C₆ alkylthio, arylthio, C₁-
20 C₆ alkoxy carbonyl, nitro, heteroarylthio, C₁-C₆
alkylsulfonyl, arylsulfonyl, and aryloxy; Sub₁₅ is selected
from the group consisting of hydrogen, C₁-C₆ alkylthio, C₁-
C₆ alkylsulfonyl, aryl, arylazo, -CH=D, cyano, C₁-C₆
alkylsulfonyl, aryl, heteroaryl, arylthio, arylsulfonyl,
25 C₁-C₆ alkoxy, trifluoromethyl and C₁-C₆ alkoxy carbonyl;
Sub₁₆ is selected from the group consisting of hydrogen,
one or two groups selected from halogen, C₁-C₆
alkylsulfonyl, nitro, cyano, C₁-C₆ alkoxy carbonyl,
thiocyano and C₁-C₆ alkylthio; Sub₁₇ is selected from the
30 group consisting of hydrogen, one or two groups selected
from C₁-C₆ alkyl, halogen, nitro and SO₂N (C₁-C₆ alkyl)₂;
Sub₁₈ is selected from the group consisting of hydrogen,
aryl, C₁-C₆ alkylthio, C₁-C₆ alkylsulfonyl and heteroaryl;
Sub₁₉ is selected from the group consisting of hydrogen,
35 one or more groups selected from C₁-C₆ alkyl, cyano, nitro,
aryl, heteroaryl, arylazo, -CH=D, C₁-C₆ alkoxy carbonyl,


09751766-122900

15






- wherein R₁₇ is selected from the group consisting of
- 5 hydrogen, 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, -C₂-C₆ alkylene-OH, O C₂-C₆ alkylene-C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene-C₁-C₆ alkanoyloxy, halogen, carboxy, C₁-C₆ alkoxycarbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄,
 - 10 NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl, R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or
 - 15 more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,




wherein R_5' , R_{16}' and Q are as defined in claim 63; R_{18} and R_{19} are independently selected from the group consisting of hydrogen, unsubstituted C_1 - C_{10} alkyl, substituted C_1 - C_{10} alkyl, C_3 - C_8 cycloalkyl, C_3 - C_8 alkenyl, C_3 - C_8 alkynyl and aryl or R_{18} and R_{19} may be combined with another element to which they are attached to form a radical Z having the formula




wherein Q_2 is selected from the group consisting of a covalent bond, $-O-$, $-S-$, $-SO_2-$, $-CO-$, $-CO_2-$, $-N-(C_1-C_6 \text{ alkyl})-$, $-N(CO \ C_1-C_6 \text{ alkyl})-$, $-N(SO_2 \ C_1-C_6 \text{ alkyl})-$, $-N(CO \text{ aryl})-$, or $-N(SO_2 \text{ aryl})-$; R_{20} , R_{21} and R_{22} are independently selected from the group consisting of or C_1 - C_6 alkyl; R_{23} is selected from the group consisting of hydrogen, C_1 - C_6 alkyl, C_3 - C_8 cycloalkyl, heteroaryl or aryl; with the provision that two carboxy ($-CO_2H$) groups be on Z , such that the two carboxy groups be present on either R_{17} or R_{18} , or one carboxy may be present on each of R_{17} and R_{18} .



91. The diacidic compounds of claim 90 wherein R_{18} is




or R_{17} is



91. The diacidic compounds of claim 89 wherein R_{18} and R_{19} are independently selected from the group consisting of




and



; or


R₁₈ is selected from the group consisting of the groups
5 listed immediately above and R₁₇ is selected from the group consisting of



10 or NHCO C₁-C₄ alkylene CO₂H; wherein Q is selected from the group consisting of -S-, -O- or -SO₂-.


92. A method comprising reacting

- a) at least one diacidic monomer, comprising
15 about 1 to 100 mole % of at least one light-absorbing monomer having a light absorption maximum between about 300 nm and about 1200 nm and 99-0 mole % of a non-light absorbing monomer which does not absorb significant light at wavelengths above 300 nm or has a light absorption
20 maximum below 300 nm, with
b) an organic compound having the formula




09751756 122900

wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene- C₃- C₈-cycloalkylene- C₁-C₄ alkylene, C₁-C₄ alkylene-arylene- C₁-C₄ alkylene, C₂- C₄-alkylene-L-
 5 arylene-L- C₂-C₄ alkylene and C₂-C₄ alkylene-(L- C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from the group consisting of -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; X and X₁ are
 10 reactive groups and are independently selected from the group consisting of bromine, iodine and R-SO₂O; wherein R is selected from the group consisting of C₁-C₆ alkyl; C₁-C₆ alkyl substituted with chlorine, fluorine, C₁-C₆ alkoxy, aryl, aryloxy, arylthio or C₃-C₈ cycloalkyl; C₃-C₈
 15 cycloalkyl or aryl, wherein said reaction is carried out in a solvent in the presence of a base to form a light absorbing composition comprising a mixture of a polymer having the formula



20

and a cyclic compound having the general formula



wherein B is as defined above, n is at least 2, m is 1, 2,
 25 3 or 4 and A comprises the residue of said diacidic monomer.

93. The process of claim 92 where said light-absorbing monomers have the formula

30



09751766-122900


wherein H represents an acidic hydrogen atom; Y is a
divalent light-absorbing moiety selected from the group
consisting of chromophoric classes of azo, disazo, triazo,
bis-azo, methine, arylidene, polymethine, azo-methine,
5 azamethine, anthraquinone, anthrapyridone (3H-
dibenz[f,ij]isoquinoline-2,7-dione, nitroarylamines
anthrapyridine (7H-dibenz[f,ij]isoquinoline-7-one,
phthaloylphenothiazine (14H-naphth[2,3-a]phenothiazine-
8,13-dione, benzanthrone (7H(de)anthracene-7-one),
10 anthrapyrimidine (7H-benzo[e]perimidine-7-one),
anthrapyrazole, anthraisothiazole, triphenodioxazine,
thiaxanthene-9-one, fluorindine (5,12-dihydroquinoxaline
[2,3-b]phenazine, quinophthalone, phthalocyanine, metal
phthalocyanine, naphthalocyanine, metal naphthalocyanine,
15 nickel dithiolenes, squarylium compounds, croconium
compounds, coumarin (2H-1-benzopyran-2-one), coumarin
imine (2H-1-benzopyran-2-imine), perinone, benzodifuran,
phthaloylacridone, phthaloylphenoxazine (14H-naphtho[2,3-
a]phenoxazine-8,13-dione, phthaloylacridone (13H-
20 naphtho[2,3-c]acridine-5,8,14-trione),
anthraquinonethioxanthene (8H-naphtho[2,3-c]thioxanthene-
5,8,13-trione, anthrapyridazone, pyrrolo[3,4-c]pyrrole,
indigo, thioindigo, quinoline, xanthene, acridine, azine,
cyanine, oxazine, 1,4 and 1,5-naphthoquinones, 2,5-
25 diarylaminoterephthalic acids and esters, pyromellitic
acid diimide, naphthalene-1,4,5,8-tetracarboxylic acid
diimide, 3,4,9,10-perylenetetracarboxylic acid diimide, 3-
aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-
oxopyrroline, arylisoindoline, hydroxybenzophenone,
30 benotriazole, naphthotriazole, diminoisoindoline,
naphthopyran (3H-naphtho[2,1-6]pyran-3-one and 3-imine,
phthalimides, 2-arylbenzazoles, carbostyryls, 1,2-
diarylethenes, 2,5-diarylthiophenes, 2,5-diaryl-1,3,4-
oxadiazoles, triazines, 2,5-diarylfurans, 2,5-diaryl-
35 1,3,4-thiadiazoles, thiophenes, 1,3-diphenyl-2-
pyrazolines, 2-arylbenzofurans, 2,6-diphenylbenzofurans,

09751766-122900

quinolines, quinoxalines, 3,4-diarylfuanones, distyrylarenes, benzanthrones, polyarenes and naphthalimides.


94. The method of claim 93 wherein said acidic functional groups are independently selected from the group consisting of $-\text{CO}_2\text{H}$, $-\text{SH}$, $-\text{OH}$ attached to an aromatic ring, $-\text{CONHCO}-$, $-\text{SO}_2-\text{NH}-\text{CO}-$, $-\text{SO}_2-\text{NH}-\text{SO}_2-$, 1(H)-1,2,4-triazol-3-yl-, imidazolyl, benzimidazolyl, pyrazolyl, $-\text{SO}_2\text{H}$ attached to aromatic ring, $-\text{NHSO}_2\text{R}_5$ and $-\text{SO}_2\text{NHR}_5$, wherein R_5 is selected from the group consisting of C_1-C_6 alkyl, C_3-C_8 cycloalkyl, aryl and C_1-C_6 alkyl substituted with at least one group selected from the group consisting of C_1-C_6 alkoxy, aryl, aryloxy, arylthio and C_3-C_8 cycloalkyl.

95. The method of claim 92 wherein said non light-absorbing monomers have the formula




wherein H represents an acidic hydrogen atom; Y_1 is a divalent moiety selected from the group consisting of $-\text{O}_2\text{C}-\text{R}_1-\text{CO}_2-$ and $-\text{O}-\text{R}_2-\text{O}-$ and $-\text{O}_2\text{C}-\text{R}_3-\text{O}-$, wherein R_1 is selected from the group consisting of C_2-C_{12} alkylene, 1-4-cyclohexylene, arylene, arylene-O-arylene, arylene- SO_2 -arylene, arylene-S-arylene, and C_1-C_4 alkylene-O- C_1-C_4 alkylene; wherein R_2 is selected from the group consisting of arylene, arylene-O-arylene, arylene-S-arylene, arylene- SO_2 -arylene, phenylene-phenylene, and phenylene- $\text{C}(\text{R}_4)_2$ -phenylene; wherein R_4 is selected from the group consisting of hydrogen and C_1-C_4 alkyl; wherein R_3 is selected from arylene.

96. A light absorbing composition comprising a mixture of a polymer having the formula



09751766.122900

and a cyclic compound having the general formula




wherein A₁ comprises the residue of at least one diacidic monomer having a light absorption maximum between about 300 nm and about 1200 nm, and wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene- C₃-C₈-cycloalkylene- C₁-C₄ alkylene, C₁-C₄ alkylene-arylene- C₁-C₄ alkylene, and C₂- C₄-alkylene-L-arylene-L- C₂-C₄ alkylene and C₂-C₄ alkylene-(L- C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; wherein n is at least 2 and m is 1, 2, 3 or 4.

97. A composition comprising a thermoplastic polymer blended with at least one light absorbing linear polymeric composition of claim 96.

98. The composition of claim 97 wherein said thermoplastic polymer is selected from the group consisting of polyesters, polyolefins, polyamides, polyimides, polyvinyl chloride, polyurethanes, polycarbonates, cellulose esters, polyacrylates, polyvinylesters, polyester-amides, polystyrene, polyacrylonitrile-butadiene- styrene, polystyrene-acrylonitrile and mixtures and blends thereof.

99. The composition of claim 96 wherein A₁ comprises the residue of at least one diacidic monomer having the structure



wherein H represents an acidic hydrogen atom and Y is a divalent light-absorbing moiety selected from the chromophoric classes consisting of azo, disazo, triazo, bis-azo, methine, arylidene, polymethine, azo-methine,

09751766-122900

azamethine, anthraquinone, anthrapyridone (3H-dibenz[f,ij]
isoquinoline-2,7-dione, nitroarylamines anthrapyridine
(7H-dibenz[f,ij]isoquinoline-7-one, phthaloylphenothiazine
(14H-naphth[2,3-a]phenothiazine-8,13-dione, benzanthrone
5 (7H-de)anthracene-7-one), anthrapyrimidine (7H-
benzo[e]perimidine-7-one), anthrapyrazole,
anthraisoithiazole, triphenodioxazine, thiaxanthene-9-one,
fluorindine (5,12-dihydroquinoxaline [2,3-b]phenazine,
quinophthalone, phthalocyanine, metal phthalocyanine,
10 naphthalocyanine, metal naphthalocyanine, nickel
dithiolenes, squarylium compounds, croconium compounds,
coumarin (2H-1-benzopyran-2-one), coumarin imine (2H-1-
benzopyran-2-imine), perinone, benzodifuran,
phthaloylacridone, phthaloylphenoxazine (14H-naphtho[2,3-
15 a]phenoxazine-8,13-dione), phthaloylacridone (13H-
naphtho[2,3-c]acridine-5,8,14-trione),
anthraquinonethioxanthane (8H-naphtho[2,3-c]thioxanthene-
5,8,13-trione, anthrapyridazone, pyrrolo[3,4-c]pyrrole,
indigo, thioindigo, quinoline, xanthene, acridine, azine,
20 cyanines, oxazine, 1,4 and 1,5-naphthoquinones, 2,5-
diarylaminoterephthalic acids and esters, pyromellitic
acid diimide, naphthalene-1,4,5,8-tetracarboxylic acid
diimide, 3,4,9,10-perylenetetracarboxylic acid diimide, 3-
aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-
25 oxopyrroline, arylisoindoline, hydroxybenzophenone,
benzotriazole, naphthotriazole, diminoisoindoline,
naphthopyran (3H-naphtho[2,1-6]pyran-3-one and 3-imine,
phthalimides, 2-arylbenzazoles, carbostyryls, 1,2-
diarylethenes, 2,5-diarylthiophenes, 2,5-diaryl-1,3,4-
30 oxadiazoles, triazines, 2,5-diarylfurans, 2,5-diaryl-
1,3,4-thiadiazoles, thiophenes, 1,3-diphenyl-2-
pyrazolines, 2-arylbenzofurans, 2,6-diphenylbenzofurans,
quinolines, quinoxalines, 3,4-diarylfuanones,
distyrylarenes, benzanthrone, polyarenes and
35 naphthalimides.

09751766-122900

100. The light absorbing composition of claim 99 wherein A₁ further comprises less than about 50% by weight of the total composition of a residue of at least one non-light absorbing monomer having the formula



wherein Y₁ is a divalent moiety, selected from the group consisting of -O₂C-R₁-CO₂- and -O-R₂-O- and -O₂C-R₃-O-, wherein R₁ is selected from the group consisting of C₂-C₁₂ alkylene, 1-4-cyclohexylene, arylene, arylene-O-arylene, arylene-SO₂-arylene, arylene-S-arylene, and C₁-C₄ alkylene-O- C₁-C₄ alkylene; wherein R₂ is selected from the group consisting of arylene, arylene-O-arylene, arylene-S-arylene, arylene-SO₂-arylene, phenylene-phenylene, and phenylene-C(R₄)₂-phenylene; wherein R₄ is selected from the group consisting of hydrogen and C₁-C₄ alkyl; wherein R₃ is arylene; wherein B is a divalent organic radical selected from the group consisting of C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene-C₃-C₈-cycloalkylene-C₁-C₄ alkylene, C₁-C₄ alkylene-arylene-C₁-C₄ alkylene, and C₂-C₄-alkylene-L-arylene-L-C₂-C₄ alkylene and C₂-C₄ alkylene-(L-C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -N(SO₂ C₁-C₆ alkyl)-, -N(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof.

101. The light absorbing composition of Claim 100 wherein A₁ comprises the residue of at least one diacidic monomer having at least one acidic group other than carboxy and having a light absorption maximum between about 300 nm and about 1200 nm and wherein B is a divalent organic radical selected from C₂-C₁₂ alkylene, C₃-C₈ cycloalkylene, C₁-C₄ alkylene- C₃- C₈-cycloalkylene- C₁-C₄ alkylene, C₁-C₄ alkylene-arylene- C₁-C₄ alkylene, and C₂-C₄-alkylene-L-arylene-L- C₂-C₄ alkylene and C₂-C₄ alkylene-(L-C₂-C₄ alkylene)₁₋₄, wherein L is a linking group selected from -O-, -S-, -SO₂-, -NH-, -N(C₁-C₆ alkyl)-, -N(aryl)-, -

09751766-122900









~~N(SO₂ C₁-C₆ alkyl)-, -(SO₂aryl)-, -SO₂N(C₁-C₆ alkyl)- and combinations thereof; wherein n is at least 2.~~

~~102. The composition of claim 99 wherein said light absorbing monomer comprises two acidic groups independently selected from the group consisting of -CO₂H, SH, hydroxy attached to an aromatic ring, -CONHCO- (imide), -SO₂NHCO-, -SO₂NHSO₂-, 1(H)-1,2,4-triazolyl-3-yl-, imidazolyl, benzimidazolyl, pyrazolyl, -SO₂H attached to an aromatic ring, -NHSO₂R₅ and -SO₂NHR₅, wherein R₅ is selected from the group consisting of C₁-C₆ alkyl; C₁-C₆ alkyl substituted with at least one group selected from C₁-C₆ alkoxy, aryl, aryloxy, arylthio and C₃-C₈ cycloalkyl; C₃-C₈ cycloalkyl; aryl.~~


~~103. The composition of claim 100 wherein said light absorbing monomer comprises two carboxy groups as acidic functional groups.~~

~~104. The composition of claim 100 wherein the light absorbing portion of A comprises the residue of at least one diacidic light absorbing monomer selected from the group consisting of the anthraquinone and anthrapyridone colorants having the structures:~~


00622T 9945460



006221 " 9975760




- wherein R_{14} is selected from the group consisting of
- hydrogen and 1-4 groups selected from amino, C_1-C_{10}
 - 5 alkylamino, C_3-C_8 alkenylamino, C_3-C_8 alkynylamino, C_3-C_8 cycloalkylamino, arylamino, halogen, C_1-C_6 alkoxy, C_1-C_6 alkylthio, aryl, aroyl, C_1-C_6 alkanoyl, C_1-C_6 alkanoyloxy, NHCO C_1-C_6 alkyl, NHCOaryl, NHCO C_1-C_6 alkyl, NHO C_1-C_6 alkyl, NHO aryl, C_1-C_6 alkoxycarbonyl, aryloxy, arylthio,
 - 10 heteroarylthio, cyano, nitro, trifluoromethyl, thiocyno, SO C_1-C_6 alkyl, SO aryl, $-SO_2NH$ C_1-C_6 alkyl, $-SO_2N(C_1-C_6$ alkyl) $_2$, $-SO_2N(C_1-C_6$ alkyl) aryl, CONH C_1-C_6 alkyl, CON(C_1-C_6 alkyl) $_2$, CON(C_1-C_6 alkyl) aryl, C_1-C_6 alkyl, furfurylamino, tetrahydrofurfurylamino, 4-(hydroxymethyl)
 - 15 cyclohexanemethylamino,



or hydroxy; Q and Q' are independently selected from the
 5 group consisting of -O-, -N(COR₁₀)-, -N(SO₂R₁₀)-, -N(R₁₀)-,
 -S-, -SO₂-, -CO₂-, -CON(R₁₀)-, SO₂N (R₁₀)-, wherein R₁₀ is
 selected from the group consisting of hydrogen, aryl, C₃-C₈
 cycloalkyl, or C₁-C₁₀ alkyl; R₁₅ is selected from the group
 10 consisting of hydrogen, cyano, C₁-C₆ alkylamino, C₁-C₆
 alkoxy, halogen, arylthio, aryl, heteroaryl,
 heteroarylthio, C₁-C₆ alkoxy carbonyl, aroyl or
 arylsulfonyl; R₁₆ is selected from the group consisting of
 hydrogen, C₁-C₆ alkyl, C₃-C₈ cycloalkyl and aryl; R₁₆' is
 15 selected from the group consisting of hydrogen, C₁-C₆
 alkyl, halogen and C₁-C₆ alkoxy; wherein each C₁-C₆ alkyl
 group and C₁-C₆ alkyl group which is a portion of another
 group may contain at least one substituent selected from
 the group consisting of hydroxy, cyano, chlorine,
 20 fluorine, C₁-C₆ alkoxy, C₃-C₈ cycloalkoxy, C₁-C₆
 alkylcyclohexyl, hydroxymethyl cyclohexyl, aryl and
 heteroaryl; with the provision that two acidic groups
 containing one acidic proton each or one acidic group
 containing two acidic hydrogens be present in the diacidic
 compounds.

25 105. The composition of claim 100 wherein the light
 absorbing portion of A comprises the residue of at least
 one light absorbing monomer selected from the group
 consisting of azo, disazo, bis-azo and azomethine and
 having respectively the structures:




wherein R₆ is the residue of an aromatic or heteroaromatic
 amine which has been diazotized and coupled with a
 35 coupling component H-Z and is derived from an amine

006221" 99475760

- selected from the group consisting of aromatic and heteroaromatic amine classes selected from the group consisting of aniline, 1-aminonaphthalene, 1-aminoanthraquinone, 4-aminoazobenzene, 2-aminothiazole, 2-aminobenzothiazole, 3-amino-2,1-benzisothiazole, 2-aminothieno[2,3-d]thiazole, 5-aminoisothiazole, 5-aminopyrazole, 4-aminopyrazoloisothiazole, 2-amino-1,3,4-thiadiazole, 5-amino-1,2,4-thiadiazole, 5-amino-1,2,3-triazole, 2-amino-1,3,4-triazole, 2(5) aminoimidazole, 3-aminopyridine, 2(3) aminothiophene, 2(3) aminobenzo[b]thiophene, 2-aminothieno[3,2-b]thiophene, 3-aminothieno[2,3-c]isothiazole, 3-amino-7-benz-2,1-isothiazole, 3-aminobenzothienoisothiazole, 3-aminoisothiazole[3,4-d]pyrimidine, 5-amino-1,2,3-triazole, 3(4) aminophthalimide and 5(6) amino-1,2-benzisothiazolon-1,1-dioxide with said aromatic and heteroaromatic ring systems being unsubstituted or substituted with one or more groups selected from C₁-C₁₀ alkyl, C₁-C₆ alkoxy, C₃-C₈ cycloalkyl, carboxy, halogen, C₁-C₆ alkoxycarbonyl, formyl, C₁-C₆ alkanoyl, C₁-C₆ alkanoyloxy, dicyanovinyl, C₃-C₈-cycloalkanoyl, thiocyno, trifluoroacetyl, cyano, carbamoyl, -CONH-C₁-C₆ alkyl, CONHaryl, CON(C₁-C₆ alkyl)₂, sulfamoyl, SO₂NH C₁-C₆ alkyl, SO₂N(C₁-C₆ alkyl)₂, SO₂NHaryl, SO₂NH C₃-C₈ cycloalkyl, CONH C₃-C₈ cycloalkyl, aryl, aroyl, -NHSO₂ C₁-C₆ alkyl, -N(C₁-C₆ alkyl)SO₂ C₁-C₆ alkyl, -NHSO₂ aryl, NHCO C₁-C₆ alkyl, NHCO C₃-C₈ cycloalkyl, NHCOaryl, NHCO₂ C₁-C₆ alkyl, NHCONH C₁-C₆ alkyl, NHCONHaryl, N(C₁-C₆ alkyl)aryl, arylazo, heteroaryl, aryloxy, arylthio, C₃-C₈ cycloalkoxy, heteroarylazo, heteroarylthio, arylsulfonyl, tricyanovinyl, aryloxysulfonyl, C₁-C₆ alkylsulfonyl, trifluoromethyl, fluorosulfonyl, trifluoromethylsulfonyl, thiocyno, hydroxy, nitro or CH=D, wherein D is the residue of an active methylene compound selected from the group consisting of malononitrile, α-cyanoacetic acid esters, malonic acid esters, α-cyanacetic acid amides, α-C₁-C₆

09751766-122900

alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles,
 α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles,
 α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-
indanediones, 2-furanones, benzo-2-furanones, naphtho-2-
5 furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-
dioxo (2H)-pyridines, benzo (b) thieno-3-ylidene propane
dinitrile-5,5-dioxides, 1,3-bis(dicyanomethylene) indanes,
barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-
dihydro-1-benzothiophene-1,1-dioxides or aryl-
10 C(CH₃)C=C(CN)₂; wherein R₇ is a divalent aromatic or
heteroaromatic radical selected from the group consisting
of 1,4-phenylene, naphthalene-1,4-diyl, thiazol-2,5-diyl
and thiophene-2,5-diyl:



15


wherein R₈ is selected from the group consisting of
hydrogen or 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆
alkoxy, cyano, halogen, -NHCO C₁-C₆ alkyl, -NHCO₂ C₁-C₆
20 alkyl, -NHCO aryl, -NHCONH aryl or NHCONH C₁-C₆ alkyl; R₉
is selected from the group consisting of hydrogen, C₁-C₆
alkyl, halogen, aryl, heteroaryl; R₁₀ is selected from the
group consisting of hydrogen, C₁-C₆ alkoxycarbonyl, cyano,
carbamoyl, aryl, arylsulfonyl, aroyl, -CONH C₁-C₆ alkyl, or
25 C₁-C₆ alkylsulfonyl; wherein Z is the residue of a coupling
compound selected from the group consisting of electron
rich compound classes of anilines, 1-aminonaphthalenes,
1,2-dihydroquinolines, 1,2,3,4-tetrahydroquinolines,
benzomorpholines (3,4-dihydro-2H-1,4-benzoxazine),

pyrazolones, pyrazoles, 3-cyano-6-hydroxy-2-pyridones,
2,3-dihydroindoles, indoles, 4-hydroxycoumarins,
4-hydroxy-2-quinolones, imidazo[2,1-b]thiazoles,
julolidines (2,3,6,7-tetrahydro-1H,5H-
5 benzo[ij]quinolizines), 1-oxajulolidines, 1,2,5,6-
tetrahydro-4H-pyrrolo[3,2,1-ij]quinolines, 2,6-diamino-3
cyanopyridines, 2-aminothiazoles, 2-aminothiophenes, 5,5-
dimethyl-1,3-cyclohexanedione (dimedone), phenols,
naphthols, 2,4-pentanediones or acetoacetarilides; wherein
10 Y₁ is the residue of a bis coupling component selected
from the group consisting of anilines, 1,2-
dihydroquinolines, 1,2,3,4-tetrahydroquinolines,
benzomorpholines (3,4-dihydro- 2H-1,4-benzoxazines), 3-
cyano-6-hydroxy-2-pyridones, 2,6-diaminopyridines, 2,3-
15 dihydroindoles, naphthylamines, 2-aminothiazoles, or a
combination of these; with the provision that two acidic
functional groups containing one acidic hydrogen each or a
functional group containing two acidic hydrogen are
present in the diacidic light absorbing monomer.

20




106. The composition of claim 100 wherein the light
absorbing portion of A comprises the residue of at least
one light absorbing monomer selected from the group
consisting of methine, arylidene, polymethine, azamethine,
25 3-aryl-2,5-dioxypyrroline, 3-aryl-5-dicyanomethylene-2-
oxypyrroline and arylisoindoline and having respectively
the structures:

00661766.1 22900




methylene compound selected from the group consisting of malononitrile, α -cyanoacetic acid esters, malonic acid esters, α -cyanacetic acid amides, α -C₁-C₆ alkylsulfonylacetonitriles, α -arylsulfonylacetonitriles, α -C₁-C₆ alkanoylacetonitriles, α -aroylacetonitriles, α -heteroarylacetonitriles, bis(heteroaryl)methanes, 1,3-indanediones, 2-furanones, benzo-2-furanones, naphtho-2-furanones, 2-indolones, 3-cyano-1,6-dihydro-4-methyl-2,6-dioxy (2H)-pyridines, benzo (b) thieno-3-ylidene propane dinitrile-5,5-dioxides, 1,3-bis(dicyanomethylene) indanes, barbituric acid, 5-pyrazolones, dimedone, 3-oxo-2,3-dihydro-1-benzothiophene-1,1-dioxides or aryl-C(CH₃)C=C(CN)₂, with the proviso that two acidic functional groups containing one acidic hydrogen each, or a functional group containing two acidic hydrogens are present.

107. The composition of claim 100 wherein the light absorbing portion of A₂ comprises a residue of at least one diacidic coumarin compounds selected from the group consisting of the structures



wherein Z₃ is selected from the group consisting of cyano, C₁-C₆ alkoxy carbonyl, C₁-C₆ alkylsulfonyl, arylsulfonyl, aryl, heteroaryl, formyl, aroyl, C₁-C₆ alkanoyl or -CH=D, wherein R₁₇ is selected from the group consisting of hydrogen, 1-2 groups selected from C₁-C₆ alkyl, C₁-C₆ alkoxy, C₁-C₆ alkylthio, -O C₂-C₆ alkylene-OH, O C₂-C₆ alkylene- C₁-C₆ alkanoyloxy, C₁-C₆ alkylene-OH, C₁-C₆ alkylene- C₁-C₆ alkanoyloxy, halogen, carboxy, C₁-C₆ alkoxy carbonyl, trifluoromethyl, NHCOR₂₄, NHCO₂R₂₄,


NHCON(R₂₄)R₂₅, and NHSO₂R₂₅, wherein R₂₄ is selected from the group consisting of hydrogen, C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl, R₂₅ is selected from the group consisting of C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl or aryl wherein each C₁-C₁₀ alkyl group in R₂₄ and R₂₅ may be further substituted with one or more groups selected from the group consisting of C₃-C₈ cycloalkyl, aryl, aryloxy, arylthio, CO₂H, CO₂ C₁-C₆ alkyl, cyano, hydroxy, succinimido, C₁-C₆ alkoxy,



10

wherein R₅' is selected from the group consisting of hydrogen, C₁-C₆ alkyl or aryl; R₁₆' is selected from hydrogen or one or two groups selected from C₁-C₆ alkyl, halogen, and C₁-C₆ alkoxy; Q is selected from the group consisting of -O-, -N(COR₁₀)-, -N(R₁₀)-, -S-, -SO₂-, -CO₂-, CON(R₁₀), SO₂(R₁₀)-, wherein R₁₀ is selected from the group consisting of hydrogen, aryl, C₃-C₈ cycloalkyl or C₁-C₁₀ alkyl; R₁₈ and R₁₉ are independently selected from the group consisting of hydrogen, unsubstituted C₁-C₁₀ alkyl, substituted C₁-C₁₀ alkyl, C₃-C₈ cycloalkyl, C₃-C₈ alkenyl, C₃-C₈ alkynyl and aryl or R₁₈ and R₁₉ may be combined with another element to which they are attached to form a radical Z having the formula

25



006221 96 132900

25

add
c27